

STIC-ILL

241056

From: Schwadron, Ronald
Sent: Wednesday, February 10, 1999 11:02 AM
To: STIC-ILL
Subject: 08902692

Youdim et al.
Clin. Ecology
vol 7
p 55-61
1990

Youdim et al.
Clin. Ecology
vol 7
p 62-66
1990

Youdim et al.
Clin. Ecology
vol 7
p 67-72
1990

Rea
Annual Meeting-American Academy of Environmental Medicine (I think this is a book of abstracts published by
American academy of Environmental Medicine)
Sept 1995
p213-222

0735-9306

Griffiths et al.
Cytobios
vol 93
p49-66
1998

thanxs

ask
2/11

VRG-NO
2/11

gto
NLm
2/19
104


NIH-NOS
2/10 RC

Archives for Human Ecology in Health & Disease

W1 CL694W
V.7 NO.3-4 1990
C.01-----SEQ: C25190000
TI: CLINICAL ECOLOGY

Clinical Ecology

Volume 7, Number 3 & 4, 1990



TREATMENT OF ENVIRONMENTALLY SENSITIVE PATIENTS WITH TRANSFER FACTOR

PART III: CASE STUDIES ON THREE PATIENTS

Said Youdim, Ph.D.¹
Hsueh-chia Liang, M.D.²
William J. Rea, M.D.³

ABSTRACT

Three case studies of environmentally sensitive patients treated with transfer factor (TF) are presented in this paper. Case 1 is a 44-year-old female with multiple allergies, hyper IgE and severe dermatitis. Case 2 is a 51-year-old female with multiple food and inhalant allergies and increased levels of blood chemicals. Case 3 is a 46-year-old female also with multiple allergies and environmental sensitivities. Upon treatment with TF, Case 1 demonstrated a most dramatic response with clearance of her eczematous skin, decrease in her serum IgE level, improved delayed cutaneous hypersensitivity (cellular immunity) and increase in the number of her suppressor cells. Case 2 demonstrated rapid clinical progress associated with improved immunologic parameters and decreased levels of toxic blood chemicals. Case 3, a highly reactive individual, showed slow but steady progress clinically and immunologically with increase in the number of lymphocytes, T cells, T cell subsets and CMI. This study indicated the diversity of patient population and response to TF therapy. *Transfer factor, T & B lymphocytes, cell mediated immunity, eczema, phenotype.*

INTRODUCTION

In previous communications (1,2), we reported on the immunological and clinical response of a group of patients placed on a course of transfer factor (TF) therapy for a period of 6-12 months. The data presented the overall picture of our findings and the accumulated results. In this communication we present three representative case reports in order to familiarize the reader with specific examples of the patient population and their response to TF.

MATERIALS AND METHODS

Technical data patient selections, questionnaires, TF preparation and doses were as described previously (1,2). Immunoglobulin E determinations were performed using the Kallestad Quanti Clone kits (Kallestad Laboratories, Austin, Texas). Chemistries were performed by local laboratories. Blood toxic chemical levels were performed by AccuChem Laboratories (Richardson, Texas). Sensitivity to foods, molds, chemicals and hormones were determined by inhalation or intradermal challenge (3,4,5,6).

CASE SUMMARIES AND RESULTS

Case #1

A 44-year-old white female was admitted to the Environmental Control Unit with the chief complaint of urticaria. The patient reported that she had eczema and asthma as a child, and was hospitalized at the age of 1 for eczema. In mid-1978, she married and developed a severe erythematous rash with spreading of her eczema to her face, neck, arms, chest, and thighs. Burning, itching "sores" were a major problem. She noted that when she returned to her home in New Orleans there was a foam pad on her

¹11927 Darlington Ave, #4
Los Angeles, CA 90049

²Research Fellow, Environmental Health Center - Dallas

³Environmental Health Center - Dallas

mattress and upon exposure she had extreme whole-body dermatitis. She reported that certain foods and exposure to detergents and several other products caused the skin condition to worsen. She also had several flares of generalized urticaria from foods. She had tried many forms of therapy, but had only responded to steroids.

Her past medical history included a tonsillectomy and adenoidectomy in 1948, a benign breast tumor was excised in 1969. She had had several skiing accidents including bone fractures. There was a familial history of hay fever, cancer, eczema, depression, asthma, psychiatric care, heart disease, drug use, arthritis, drug allergies, and diabetes. Her mother died at the age of 54 of breast cancer. Her father was aged 70 with known salicylate problems, and sinus troubles. Two brothers both suffered from hay fever, and a sister had eczema.

Physical examination was normal except for the skin which showed scaly, erythematous rash over the upper trunk and extremities and upper legs. Rough, scaly skin all over the body was noted. Many areas were moist with exudate.

Intradermal tests showed positive reactions to 50 foods, molds, lake algae, T.O.E., Candida, house dust mite, smut, histamine, serotonin II, trees, weeds, terpenes, flagon, MRV, Nystatin, orris root, newsprint, perfume, formaldehyde and phenol.

Chemical challenges to assess her sensitivities were done in a steel and glass airtight booth by double blind method (3). Chemicals that produced symptoms were phenol (<.002ppm), an insecticide (<.034ppm), chlorine (<.33ppm), petroleum derived ethanol (<.5ppm) and normal saline.

She was asked to keep an environmentally clean oasis at home, eat less chemically contaminated foods and water, and take antigen injections for foods and inhalants. This resulted in some improvement, allowing the cessation of steroids. She found that heat made her more ill, so she moved from New Orleans to Chicago with benefit. The patient also found that with food immunotherapy her adverse foods reactions were controlled. She also noted that her symptoms were worse in the country, when exposed to plants, pollens and terpenes.

The patient was placed on 2 units of transfer factor per week on 10/23/86. This resulted in a diminution in her allergies, fatigue and an improvement in her ability to concentrate (see patient questionnaire refs. 1, 2). Her skin began to clear and her lymphocytes, total T cells, T helper and T suppressor cells increased significantly (Table 1). Her cell mediated immunity improved markedly (Table 2) with the positive response rising from one to four antigens and the sum of the mean reaction size growing from 2 to 26mm. At the same time there was a dramatic decrease in her IgE level (Table 1).

TABLE 1
Sequential WBC, T Cells Numbers and IgE Levels for Case 1

Immune Parameters			Before Treatment		After Treatment		Normal Range
	1981	1982	1983	1984	1985	1988	
WBC	6800	8700	5800	7000	7100	7500	4800-10000/mm ³
Lymphocyte(%)	0017	0021	0021	----	0014	0027	21-49
Total Lymphs	1156	1827	1218	1470	0994	2025	1400-4200/mm ³
T11 (%)	0073	0050	0065	0071	0082	0085	62-86
Total T11	0844	0914	0792	1044	0815	1721	1260-2560/mm ³
T4 (%)	----	----	----	0074	0060	0071	32-54
Total T4	----	----	----	1088	0596	1438	0670-1800/mm ³
T8 (%)	----	----	----	0008	0021	0012	17-35
Total T8	----	----	----	0118	0209	0243	0333-1070/mm ³
T4/T8	----	----	----	0009.25	0003.0	0005.9	01-2.7
B (%)	0022	----	0010	----	0013	0010	05-18
Total B	0254	----	0122	----	0129	0203	0082-0411/mm ³
Eosinophils (%)	0037	0011	0017	----	0012	0001	0-4%
Eos. Total	2182	0968	1197	----	----	----	0150-0250/mm ³
IgE	1048	0610	>8000	----	18400	13156	10-15 IU/ml

TABLE 2
Increase in CMI After Transfer Factor Therapy, Case 1

Antigens Tested	Before Treatment 1985	After Treatment 1987
Tetanus	2	0
Diphtheria	0	7
Streptococcus	0	8
Tuberculin	0	5
Control	0	0
Candida	0	6
Trichophyton	0	0
Proteus	0	0
Scoring		
Number of positive reactions	1	4
Sum of average (mm)	2	26

Case #2

This 51-year-old white female was admitted with the chief complaints of urinary frequency and urgency, abdominal bloating, cramping, muscle and joint pain and

headaches. She had had ongoing symptoms since childhood with numerous respiratory infections. Additionally during high school she developed chronic constipation which has persisted to date. Diagnostic evaluation by intradermal challenge showed the patient was sensitive to 22 foods, 4 weeds, 4 trees, 3 grasses, 6 chemicals, hormones, 7 molds, 2 smuts and histamine. The patient was treated with immunotherapy for her sensitivities. She also received supportive therapy with nutritional, vitamin and mineral supplementation.

She was placed on two units of transfer factor per week in 1987. Ten months after the start of TF therapy, she reported improvements in her allergic reactions, recurrent infections, fatigue, concentration, bowel problems and depression (see patient questionnaire 1, 2). Table 3 shows the patient's WBC and lymphocyte counts before and after seven months of TF therapy. There was a marked increase in her T cells and T cell sub populations. Her CMI response (Table 4) also improved from one barely positive reaction of 3mm to two reactions with a sum of 11mm in 1988 and to 6 reactions of 30mm in 1989. Her response to diphtheria antigen increased, and she developed de novo responses to strep, tuberculin, candida, trichophyton and proteus.

TABLE 3
WBC, Lymphocyte and T Cell Numbers for Case 2

Immune Parameters	Before Treatment 10/20/87	After Treatment 5/19/88	Normal Range
WBC	5200	4900	4800-10000/mm ³
Total lymphs	0728	1470	1400-4200
Lymphocyte %	0014	0030	0021-0049
T11 %	0089	0088	0062-0086
Total T11	0648	1294	1260-2650/mm ³
T4 %	0068	0058	0032-0054
Total T4	0495	0853	0670-1800/mm ³
T8 %	0014	0016	0017-0035
Total T8	0102	0235	0333-1070/mm ³
T4/T8	0004.8	0003.6	0001-0002.7
B %	0010	0003	0005-0018
Total B	0073	0044	0082-0477/mm ³

the patient's WBC and lymphocyte counts before and after seven months of TF therapy. There was a marked increase in her T cells and T cell sub populations. Her CMI response (Table 4) also improved from one barely positive reaction of 3mm to two reactions with a sum of 11mm in 1988 and to 6 reactions of 30mm in 1989. Her response to diphtheria antigen increased, and she developed de novo responses to strep, tuberculin, candida, trichophyton and proteus.

Case #3

This 46-year-old white female was admitted with the chief complaint of right frontal sinus headaches for the past 9 years. Other symptoms were recurrent nausea and vomiting, gas, abdominal bloating and occasional sore throat. Recurrent fatigue and lassitude was a large problem as well. Additionally, she had adrenal insufficiency and had also had several operations in 1975 and 1976. The

TABLE 4
Increase in CMI After Transfer Factor Therapy, Case 2

Antigens Tested	Diameter in Millimeters		
	Before Treatment 1987	After Treatment	
		1988	1989
Tetanus	0	0	0
Diphtheria	3	6	4
Streptococcus	0	0	4
Tuberculin	0	0	5
Control	0	0	0
Candida	0	0	5
Trichophyton	0	0	4
Proteus	0	5	6
Scoring			
Number of positive reactions	1	2	6
Sum of average (mm)	3	11	30

TABLE 5
WBC, Lymphocyte and T Cell Numbers for Case 3

Immune Parameters	Before Treatment		After Treatment	Normal Range
	9/2/85	4/1/86	5/6/87	
WBC	5100	4000	4600	4800-10000/mm ³
Lymphocytes %	0023	0030	0058	0021-0049
Total Lymphocytes	1173	1200	2669	1600-4200/mm ³
T11	0072	0080	0087	0062-0086
Total T11	0845	0960	2231	1260-2650/mm ³
T4 %	0054	0044	0048	0032-0054
Total T4	0633	0528	1281	0670-1800/mm ³
T8 %	0019	0034	0025	0017-0035
Total T8	0223	0408	0667	0333-1070/mm ³
T4/T8	0002.8	0001.3	0001.9	0001-0002.7
B Lymphocyte %	0005	0008	0005	0005-0018
Total B	0059	0096	0133	0082-0477/mm ³

following food challenges produced symptoms of headaches, fatigue, nausea and vomiting: pork, turkey, pineapple, rabbit, perch, broccoli, frog legs, duck, millet, moose, spinach, coconut, venison, catfish, goat milk, strawberries, barley, lima beans, dates, grapefruit, beef, eggs and zucchini.

Inhalation of phenol (0.4%) for less than 10 minutes produced yawning, sleepiness, nausea, confusion and fatigue. Natural gas exposure for 21 minutes caused tightness of sinuses, hands tingling, heart pounding, shortness of breath, dizziness, nausea and pressure headache. Inhalation of ethanol (50%) for less than 5 minutes triggered nasal stuffiness, sleepiness, yawning and postnasal discharge. Three sniffs of insecticide (Raid liquid ant and roach killer) caused nasal stuffiness, a cough, yawning and watery eyes. When exposed to formaldehyde for less than 10 minutes, she developed nasal congestion, itchy eyes, nausea, tight sinuses and sleepiness. Five minutes of challenge with chlorine (5.2%) caused nasal congestion, sore throat, watery eyes, and yawning.

The patient received antigen immunotherapy and was placed on two units of transfer factor per week in 1985. Progress was slow, but she reported a diminution in her allergic reactions, headache, recurrent infections and fatigue. (See patient questionnaire refs. 1, 2). Table 5 shows the patient's WBC and lymphocyte enumeration before and after TF therapy. Data gathered in 1987 showed a marked increase in her total lymphocyte count, total T cells and T cell populations. Table 6 shows the patient's CMI response in 1986 and 1987. Both these tests were obtained after initiation of TF therapy. CMI reactions prior to TF therapy were not available, nevertheless definite improvements in both the number and the size of the reactions was noted as therapy progressed.

TABLE 6
Increase in CMI During Transfer Factor Therapy
Case 3

Antigen Tested	Diameter (mm)	
	1986	1987
Tetanus	0	0.0
Diphtheria	4	0.5
Streptococcus	0	0.0
Tuberculin	9	5.5
Control	0	3.0
Candida	0	4.5
Trichophyton	0	3.0
Proteus	0	4.0
# of Positive Reactions	2.0	5.0
Sum of Average (mm)	13	25

DISCUSSION

Patient M.S. (Case 1) is a 44 year old female with multiple allergies, hyper IgE and severe dermatitis. Despite years of immunotherapy, this patient made modest progress until placed on TF therapy. The most dramatic response was the clearing of her eczema, the decrease in her IgE level and the increase in the number of her suppressor cells. An association between numerical deficiency of immunoregulatory T suppressor cells and atopic eczema has been reported (7,8). Abnormalities in polymorphonuclear and mononuclear cell chemotaxis and in cell mediated immunity have also been described in hyper IgE syndromes (9). Jones, et. al. (10) reported the use of oral bovine TF (OTF) in a child with skin eruptions due to hypersensitivity to eggs and milk with repeated skin infections, oral thrush and chronic cough. Administration of TF twice weekly for 6 months led to clinical improvements with associated laboratory data. These results closely parallel our findings and indicate possible usefulness of TF in these patients.

M.H. (Case 2) is a more typical patient in that she had been seen by numerous physicians including a neurologist, gynecologists, family practitioners, chiropractors and allergists without much improvement in her symptoms. She is also of interest as the improvement in her symptoms was rapid and was associated with a substantial decrease in the levels of the toluene, trichloroethylene and heptachlor epoxide in her blood. Although the association between TF therapy and a fall in the levels of toxic chemicals in the blood is not the purpose of this report, we have other data which supports this possibility. This patient also had a substantial increase in the number of her lymphocytes, T cells and T cell subsets.

Patient V.B. (Case 3) is a very sensitive patient with multiple allergies and sensitivities of long duration. Her progress was slow, but nevertheless dramatic, both clinically and as determined by laboratory data (Tables 5, 6). The number of her T cells as well as her CMI responses increased substantially during TF therapy. She was one of very few patients who reacted to the control glycerin on the CMI test (Table 6).

The above case summaries and data provided in the accompanying communications (1,2) represent comprehensive information gathered on a set of environmentally ill patients treated with TF. We feel these studies demonstrate the efficacy, safety and usefulness of TF in the treatment in certain environmentally ill patients. Similarly Bagnato and co-workers (11) reported that the degree of bronchial responsiveness to methacholine was significantly reduced at 90 days during treatment with thymomodulin and remained reduced, even if not significantly, 60 days after cessation of treatment.

Thymomodulin is a calf thymus acid lysate capable of improving the clinical symptoms of children with atopic dermatitis (12) and food allergy (13). Further, thymomodulin has been shown to regulate maturation of human and murine Pre-T-lymphocytes, and to modulate the function of mature human T & B lymphocytes for the regulation of immunoglobulin production (14,15). The effects of thymomodulin (16) and TF (1,2,17) might well be in modifying the entire immune response. It compares with the rather specific stimulation induced by allergen injections. It may potentiate the results of immunotherapy.

REFERENCES

1. Youdim S, Rea WJ, Liang HC. Treatment of Environmentally Sensitive Patients with Transfer Factor, I. Immunologic Studies. *Clinical Ecology* 1990;7:51-61.
2. Youdim S, Rea WJ. Treatment of Environmentally Sensitive Patients with Transfer, II. Clinical Studies and Immunological Correlates. *Clinical Ecology* 1990;7:62-66.
3. Rea WJ, Ross H, Johnson AR, et al. Confirmation of Chemical Sensitivity. *Clinical Ecology* 1989;6:113-118.
4. Miller JB. Food Allergy, Provocative Testing and Injection Therapy. Springfield, Thomas, 1972.
5. King PK, Rubin WA, Fadal RG, et al. Provocation-Neutralization: A Two-Part Study - Part I. The Intracutaneous Provocative Food Test: A Multi-Center Comparison Study. *Otolaryngol Head Neck Surg* 1988;99:263-271.
6. King PK, Fadal RG, Ward WA, et al. Provocation-Neutralization: A Two-Part Study - Part II. Subcutaneous Neutralization Therapy: A Multi-Center Study. *Otolaryngol Head Neck Surg* 1988;99:272-277.
7. Leung DYM, Rhodes AR, Geha RS. Enumeration of T Cell Subsets in Atopic Dermatitis Using Monoclonal Antibodies. *J All Clin Immunol* 1981;67:450-455.
8. Zachary CB, MacDonald DM. Quantative Analysis of T-Lymphocyte Subsets in Atopic Eczema Using Monoclonal Antibodies and Flow Cytometry. *Br J Derm* 1983;108:411-422.
9. Buckley RH. Disorders of the IgE System. In: Immunologic Disorders of Infants. Stiehm R. and Fulginiti V. (Eds). Philadelphia, Saunders. 1980:274-285.
10. Jones JF, Jeter WS, Hicks MJ. Oral Transfer Factor (OTF) Use in the Hyper-IgE Syndrome. In: Immunobiology of Transfer Factor. Kirkpatrick CH, Burger DR, Lawrence HS, (Eds). New York, Academic Press. 1983:261-269.
11. Bagnato A, Brovedani P, Comina C, et al. Long-term Treatment with Thymomodulin Reduces Airway Hyperresponsiveness to Methacholine. *Ann Allergy* 1989;62:425-428.
12. Fiocchi A, Grasso U, Rottoli A, et al. A Double Blind Clinical Trial on the Effectiveness of a Thymic Derivative (Thymomodulin) in the Treatment of Children with Atopic Dermatitis. *Int J Immunotherapy* 1986;3:279-284.
13. Genove R, Guerra A. Thymomodulin in Management of Food Allergy in Children. *Int J Tissue React* 1986;8:239-242.
14. Cazzola P, Mazzanti P, Kouttab M. Update and Future Perspectives of Thymic Biological Response Modifier (Thymomodulin). *J Immunopharmacol* 1987;9:195-216.
15. Twomey JJ, Kouttab NM. Selected Phenotypic Induction of Null Lymphocytes from Mice with Thymic and Nonthymic Agents. *Cell Immunol* 1982;72:186-194.
16. Valesini G, Barnaba V, Leviero M, et al. Clinical Improvement and Partial Correction of the T Cell Defects of Acquired Immunodeficiency Syndrome (AIDS) and Lymphadenopathy Syndrome (LAS) by a Calf Thymus Acid Lysate. *Eur J Cancer Clin Oncol* 1986;4:531-532.
17. Caney JT, Lederman MM, Toossi Z, et al. Augmentation of Skin Test Reactivity and Lymphocyte Blastogenesis in Patients with AIDS Treated with Transfer Factor. *JAMA* 1987;257:651-655.

(Item 1 from file: 34)
DIALOG(R)File 34:SciSearch(R) Cited Ref Sci
(c) 1999 Inst for Sci Info. All rts. reserv.

07717369 Genuine Article#: 200PB Number of References: 207
Title: Functional somatic syndromes
Author(s): Barsky AJ (REPRINT) ; Borus JF
Corporate Source: BRIGHAM & WOMENS HOSP, DIV PSYCHIAT, 75 FRANCIS
ST/BOSTON//MA/02115 (REPRINT)
Journal: ANNALS OF INTERNAL MEDICINE, 1999, V130, N11 (JUN 1), P
910-921

ISSN: 0003-4819 Publication date: 19990601
Publisher: AMER COLL PHYSICIANS, INDEPENDENCE MALL WEST 6TH AND RACE ST,
PHILADELPHIA, PA 19106-1572

Language: English Document Type: REVIEW
Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences; CC CLIN--Current
Contents, Clinical Medicine;

Journal Subject Category: MEDICINE, GENERAL & INTERNAL

Abstract: The term functional somatic syndrome has been applied to several related syndromes characterized more by symptoms, suffering, and disability than by consistently demonstrable tissue abnormality. These syndromes include multiple **chemical sensitivity**, the sick building syndrome, repetition stress injury, the side effects of silicone breast implants, the Gulf War syndrome, chronic whiplash, the chronic fatigue syndrome, the irritable bowel syndrome, and fibromyalgia. Patients with functional somatic syndromes have explicit and highly elaborated self-diagnoses, and their symptoms are often refractory to reassurance, explanation, and standard treatment of symptoms. They share similar phenomenologies, high rates of co-occurrence, similar epidemiologic characteristics, and higher-than-expected prevalences of psychiatric comorbidity. Although discrete pathophysiologic causes may ultimately be found in some patients with functional somatic syndromes, the suffering of these patients is exacerbated by a self-perpetuating, self-validating cycle in which common, endemic, somatic symptoms are incorrectly attributed to serious abnormality, reinforcing the patient's belief that he or she has a serious disease. Four psychosocial factors propel this cycle of symptom amplification: the belief that one has a serious disease; the expectation that one's condition is likely to worsen; the "sick role," including the effects of litigation and compensation; and the alarming portrayal of the condition as catastrophic and disabling. The climate surrounding functional somatic syndromes includes sensationalized media coverage, profound suspicion of medical expertise and physicians, the mobilization of parties with a vested self-interest in the status of functional somatic syndromes, litigation, and a clinical approach that overemphasizes the biomedical and ignores psychosocial factors. All of these influences exacerbate and perpetuate the somatic distress of patients with functional somatic syndromes, heighten their fears and pessimistic expectations, prolong their disability, and reinforce their sick role. A six-step strategy for helping patients with functional somatic syndromes is presented here.

Identifiers--KeyWord Plus(R): CHRONIC-FATIGUE-SYNDROME;
IRRITABLE-BOWEL-SYNDROME; RANDOMIZED CONTROLLED TRIAL; TERM FOLLOW-UP;
MULTIPLE CHEMICAL SENSITIVITIES; POSTTRAUMATIC-STRESS-DISORDER;
COGNITIVE-BEHAVIORAL THERAPY; MASS PSYCHOGENIC ILLNESS; CARE-SEEKING
BEHAVIOR; UPPER LIMB PAIN

Cited References:

*I MED, 1996, HLTH CONS SERV PERS

(Item 38 from file: 5)
DIALOG(R)File 5: Biosis Previews(R)
(c) 1999 BIOSIS. All rts. reserv.

03444499 BIOSIS NO.: 000023017587
CONTROLLED STUDIES ON FOOD SENSITIVITY USING INJECTION THERAPY

AUTHOR: REA W J; WILLIAMS M L; SPRAGUE D E
AUTHOR ADDRESS: DALLAS, TEXAS.

JOURNAL: 37TH ACA (AMERICAN COLLEGE OF ALLERGISTS) SCIENTIFIC CONGRESS,
WASHINGTON, D.C., USA, APRIL 4-8, 1981. ANN ALLERGY 47 (2). 1981. 139.
CODEN: ANAEA
DOCUMENT TYPE: Meeting
RECORD TYPE: Citation
LANGUAGE: ENGLISH

DESCRIPTORS: ABSTRACT HUMAN RECURRENT LARYNGEAL EDEMA
CONCEPT CODES:

13216 Nutrition-Pathogenic Diets
16006 Respiratory System-Pathology
22018 Pharmacology-Immunological Processes and Allergy
34508 Immunology and Immunochemistry-Immunopathology, Tissue Immunology
35500 Allergy
00520 General Biology-Symposia, Transactions and Proceedings of
Conferences, Congresses, Review Annuals
10060 Biochemical Studies-General
12508 Pathology, General and Miscellaneous-Inflammation and
Inflammatory Disease
12512 Pathology, General and Miscellaneous-Therapy (1971-)
14006 Digestive System-Pathology
15010 Blood, Blood-Forming Organs and Body Fluids-Other Body Fluids
22502 Toxicology-Foods, Food Residues, Additives and Preservatives

BIOSYSTEMATIC CODES:

86215 Hominidae

BIOSYSTEMATIC CLASSIFICATION (SUPER TAXA):

Animals
Chordates
Vertebrates
Mammals
Primates
Humans

4/9/80 (Item 2 from file: 65)
DIALOG(R)File 65: Inside Conferences 1993-1999/May W5
(c) 1999 BLDSC all rts. reserv. All rts. reserv.

01474979 INSIDE CONFERENCE ITEM ID: CN014639864
The Potential Uses for Autogenous Lymphocyte Factors (ALF) in
Environmental Medicine

Rea, W.

CONFERENCE: American Academy of Environmental Medicine: The cutting edge
of environmental medicine-Annual meeting; 30th
ANNUAL MEETING- AMERICAN ACADEMY OF ENVIRONMENTAL MEDICINE, 1995; 30th
P: 213-222

The Academy, 1995

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts, presentations,
papers and programme



CONFERENCE SPONSOR: American Academy of Environmental Medicine
CONFERENCE LOCATION: Tucson, AZ
CONFERENCE DATE: Sep 1995 (19950) (19950)

BRITISH LIBRARY ITEM LOCATION: 1087.378000
DESCRIPTORS: environmental medicine; medicine; AAEM

DER

The ORDER command may be used for two different document ordering features: DIALOG SourceOne, providing fast copies of original patent documents and selected business articles, and DIALORDER, Dialog's Online document ordering service for requesting the complete text of documents cited in DIALOG databases or identified through other sources.

Documents available through DIALOG SourceOne include:

- U.S. patents, from 1977 forward, updated weekly.
- International patents: European patent applications from 1978 forward and granted patents from 1985 forward, updated weekly, as well as PCT international applications from 1990 forward, updated every 2 months.

To place an order, simply type the word ORDER from any Dialog file, and respond to the specific prompts for information. To order a patent, all you need is the patent number and the country of the patent.

PRICING DETAILS

for SourceOne can be found online by typing HELP SOURCE1.

DIALORDER

DIALORDER allows you to place orders for copies of documents and have them mailed directly to your Dialog mailing address, or to an alternate address you specify. You can choose from numerous document suppliers to fill your DIALORDER request. Details about each supplier, including the supplier acronym to be used in your ORDER request, are provided on the DIALOG Yellowsheets. A list of supplier names and data sheet acronyms is available online by entering HELP SUPPLRS. The individual supplier data sheets can be obtained online by entering HELP and the letter O followed by the acronym of the data sheet, e.g., HELP OINFO for the INFO data sheet.

To order items identified during an online search:

After getting your search results, enter the ORDER command followed by the appropriate set number, a slash (/), item numbers for the documents you are requesting, a space, and the supplier acronym of your choice, i.e., ORDER S3/1-5,7 INFO. Special instructions may be entered after the acronym, i.e., ORDER S7/1-3 SRC1 SEND ATTENTION BOB FALL. The complete ORDER command may contain up to 240 characters.

To order items identified during a previous Dialog search:

BEGIN the original file again and use the KEEP command with the appropriate Dialog accession numbers to store the documents in set S0, e.g., KEEP 5647728. Then simply enter the ORDER command followed by the supplier acronym, e.g. ORDER SRC1. In OneSearch, the KEEP command must be followed by the word FROM and the database number the record is from, e.g., KEEP 5645678 from 16.

To order items not previously located online:

If you have not found an online record for the document you wish to order, use the ORDER ITEM command, followed by the supplier acronym and a description of the document, e.g. ORDER ITEM DIALOG THREE DATABASE CATALOGS. The complete ORDER command may contain up to 240 characters.

Reviewing Orders

Each order that you place is assigned an order number. To review your orders, enter the command ORDER LIST (for a summary of orders placed within the last 30 days) or ORDER LIST NEW (for a summary of orders that have not yet been transmitted to the supplier). To review the text of a specific order, enter the ORDER REVIEW command followed by the order number, e.g. ORDER REVIEW RA028.

Cancelling Orders

If an order request has not yet been transmitted to the supplier, you can cancel it with the ORDER CANCEL command, e.g. ORDER CANCEL RA028. ORDER CANCEL can be abbreviated ORDER- or O-.

Additional documentation on the DIALORDER service can be obtained by calling Dialog at 800-334-2564, or 415-858-3810.

240, 909

STIC-ILL

From: Schwadron, Ronald
Sent: Wednesday, February 10, 1999 11:02 AM
To: STIC-ILL
Subject: 08902692

Youdim et al.
Clin. Ecology
vol 7
p 55-61
1990

Youdim et al.
Clin. Ecology
vol 7
p 62-66
1990

99042172449

Youdim et al.
Clin. Ecology
vol 7
p 67-72
1990

Rea
Annual Meeting-American Academy of Environmental Medicine (I think this is a book of abstracts published by
American academy of Environmental Medicine)
Sept 1995
p213-222

Griffiths et al.
Cytobios
vol 93
p49-66
1998

2/10
NO

thanxs

CDmp

The role of the T lymphocytic cell cycle and an autogenous lymphocytic factor in clinical medicine

Bertie B. Griffiths*, William J. Rea, Bradley Griffiths and Y. Pan

EHC-D Analytical Laboratory, Environmental Health Center - Dallas, 8345 Walnut Hill Lane, Suite 240, Dallas, Texas 75231, U.S.A. (*Reprint address)

Key words: cell cycle, flow cytometry, T lymphocytes, autogenous lymphocytic factor, chemical sensitivity

Abstract

In this study 315 individuals (25 controls, 290 chemically sensitive immunocompromised patients) were investigated. Each patient had been on a standard therapy of avoidance of pollutants, nutritional supplementation, and injections of antigens for foods, and biological inhalants, but did not attain their immunological competence. Peripheral lymphocytes were collected and DNA histograms were constructed. The flow cytometer was used to evaluate the cell cycle, haematological, and other immunological profiles. From the other portion of the blood specimen, lymphocytes were propagated *in vitro*, harvested, and a lysate, termed the autogenous lymphocytic factor (ALF), was prepared. When treated with ALF, 88% of these individuals showed a significant ($p < 0.001$) clinical improvement which correlated with laboratory findings, involving regulation of abnormal cell cycles, increase in total lymphocytes and subsets T_4 , T_8 , ($p < 0.05$) and cell mediated immunity (CMI) response ($p < 0.001$). The ALF presumably acts as a biological response modifier. The cell cycle and ALF provide clinical tools for diagnosis and regulation of immunological incompetence.

Introduction

The cell cycle is the ordered and orderly events of biochemical and morphological sequences, leading from the formation of a daughter cell as a result of mitosis, to the completion of the processes required for its own division into two daughter cells (Lee and Dang, 1995). It is fundamental that following mitosis, two daughter cells are produced. These cells may initiate a new cycle (G_1 phase), some cells may become nonproliferative (G_0 phase) or progress to a restriction point where they are committed to the synthesis of cellular components, (the S phase) and finally complete the cycle (the G_2M phase). The cell cycle then comprises the sum of the growth phases of a specific cell cycle. This cycle is repeated by continuously dividing cells. Even within the same organism the specific cell cycle will vary with different classes of cells. The total time that is necessary to complete the S and G_2 phases is generally constant in different cell types. It seems reasonable then, to assume that most of the time variation takes place in the G_1 phase.

To establish homeostasis, it is imperative that the cell cycle be regulated. To this end, there are biological parameters which may be employed to ascertain the normality of a particular cell cycle. The DNA content in the nucleus of a cell (2 N or diploid amount) is constant in all normal organisms (Givan, 1992). There are only two exceptions where the DNA content is not constant; the amount of DNA will vary in cells which undergo meiosis in preparation for sexual reproduction, thus containing the 1 N or haploid amount of DNA. The other exception applies to those cells which undergo DNA synthesis in preparation for mitosis. These will contain between 2 N and 4 N amounts of DNA.

By employing the concept of DNA constancy in a particular organism, and the application of flow cytometric techniques, DNA flow histograms can be constructed depicting a normal cell cycle and the identification of abnormal cycles.

The regulation of the cell cycle in eukaryotes seems to take place at two main transition points, prior to DNA replication at a point in the G₁ phase, termed the restriction point, and prior to cytokinesis at the G₂M phase boundary (Nurse and Bissett, 1981). The progression of the cell cycle from one phase to the other is mediated by specialized proteins, cyclins, and the activation of enzymes called cyclin-dependent kinases (CDKs) and by a number of positive and negative feedback loops. To date seven CDKs have been identified and designated CDK 2 to 8 (Meyerson *et al.*, 1992; Lorinez and Reed, 1984; Nurse, 1994; Sherr, 1994; Helchman and Roberts, 1994; King *et al.*, 1994; Hunter and Pines, 1994). Each CDK acts at different stages of the cell cycle and is differentially regulated by different cyclins (Meyerson and Harlow, 1994). The critical roles played by cyclins in the regulation of eukaryotic cell cycle is amply documented.

These CDKs play very significant roles in the G₁S and G₂M transitions during the mammalian cell cycle. They regulate by phosphorylation, a number of key substrates which subsequently activate a transition from G₁ to S and from G₂ to M (Kamp *et al.*, 1994). The catalytic subunits alone of these CDKs are not active and require the influence of positive regulatory subunits to ensure biochemically active protein kinase holoenzymes (Desai *et al.*, 1992). The positive regulatory subunits employed to this end are cyclins. The activity of CDKs is, therefore, regulated by both cyclins and specific phosphorylation and dephosphorylation (Gold and Nurse, 1989; Draetta and Beach, 1988; Solomon *et al.*, 1990; Gutier *et al.*, 1991; Murray and Krischner, 1989).

Cy
invert
are d
(Evan
these
amin
and i
in ye
Sherr

Th
exhib
Fang
O'Fa
while
cells
Glot
M ph
1991
ciate
cells
Hadv

In
are i
How
aspe
obse
chen
is re
supp

Ini
phoc
were
auto
logic
imm

TI
of D
chen
shot'
indiv
infor
patie

Cyclins were identified originally as proteins in the murine invertebrate cells. The concentration of these cyclins accumulate and are destroyed periodically at defined points during the cell cycle (Evans *et al.*, 1983). At present, eight cyclins have been identified; these are designated A, B, C, D, E, F, G₁, and H based on their amino acid sequences (Pines and Hunter, 1989; Xiong *et al.*, 1991), and in some instances, on genetic complementation experiments in yeast (Nurse, 1990; Murray and Krischner, 1989; Lewin, 1990; Sherr, 1993).

The influences of the cyclins are expressed differently; *e.g.* cyclin A exhibits its influence through the S-M phase (Wang *et al.*, 1990; Fang and Newport, 1991; Walker and Maller, 1991; Lehner and O'Farrell, 1990; Pagano *et al.*, 1992; Zindy *et al.*, 1992) of most cells, while cyclin B (which is conserved from yeasts to humans) propels cells into mitosis (Nurse and Bissett, 1981; Evans *et al.*, 1983; Glotzer *et al.*, 1991). Both the A and B cyclins are degraded at the M phase by ubiquitin (Ub)-dependent proteolysis (Glotzer *et al.*, 1991). G₁ cyclins (CIN 1, CIN 2, CIN 3) are assumed to associate with a p34 CDC2 homologue, p34 CDC 28, driving yeast cells into S phase (Reed, 1992; Cross, 1988; Richardson *et al.*, 1989; Hadwiger *et al.*, 1989; Wittenberg *et al.*, 1990; Nasmyth, 1990).

In the Environmental Health Center - Dallas (EHC-D), clinical cases are investigated varying from surgical to environmental illnesses. However, cases involving chemical sensitivity entail the principal aspect of our practice. Over a period of 22 years our laboratory observations show that characteristically, a percentage of the chemically sensitive individuals portray a secondary deficit which is related to immune dysfunction, principally, depression of the suppressor T-cell (Rea *et al.*, 1986, 1987).

Initial profiles of lymphocyte cell cycle progression, T and B lymphocytes, cell mediated immunity and haematograms of patients were compared with those taken subsequent to treatment with autogenous lymphocytic factor (ALF). Acting presumably as a biological response modifier, ALF was observed to modulate deregulated immune profiles.

The focus of this investigation was firstly to observe the patterns of DNA histograms of the T lymphocyte cell cycles, especially in chemically sensitive individuals. The histograms provide a 'snapshot' of the individual's present cell cycle. By comparing an individual's cell cycle with the classic normal cycle, very crucial information will be obtained to enable a scientific regulation of a patient's cell cycle. Secondly the work was to prepare a lysate

from the *in vitro* propagated peripheral T lymphocytes of an individual with an irregular T lymphocytic cell cycle, and observe the regulatory effect of this lysate, when injected subcutaneously into such an individual. Thirdly, the intention was to establish a basis for the regulation of an individual's T lymphocytes which were observed to be irregular due to varied incitants, thus restoring normal T lymphocyte functions, and the enabling of a compromised individual to cope with multiple insults to his/her immune system. Fourthly, we examined the responsiveness of T lymphocytes to autogenous lymphocytic factor (ALF) as measured by the cell mediated immunity (CMI) test, and to establish on an individual basis an index of T lymphocyte ability to respond.

From our search of databases, and to the best of our knowledge there is no documentation pertaining to the regulation of the T lymphocytic cell cycle by lysates of an individual's T lymphocytes, and its application to clinical medicine.

Materials and methods

Isolation of lymphocytes

Venous blood was collected from 315 individuals in heparinized tubes. Of these 25 were healthy employees of EHC-D; these represented normal controls. The other individuals were chemically sensitive, chronically ill patients of the EHC-D and referrals from outside physicians. Erythrocytes were separated by a modified gradient technique using heparinized CPT vacutainers (Becton Dickinson Co.). This preparation was centrifuged for 30 min at 500 x g. The lymphocyte layer (interphase between isolymp and plasma) was removed to sterile polystyrene tubes. Lymphocytic layers were washed three times with normal saline (0.9% sodium chloride) by centrifugation in a refrigerated centrifuge at 500 x g for 20 min. Cells were resuspended in normal saline and counted.

Lymphocyte phenotyping

Monoclonal antibodies against pan T cells, T₁₁ (CD2), T helper cells T₄ (CD4), T suppressor/cytotoxic cells T₈ (CD8), activated T cells TA1 (CD26) and pan B cells B₄ (CD19) were bought from Coulter Immunology (Hialeah, Florida). Peripheral lymphocytes were obtained by venipuncture, and stained by the use of Coulter Q prep Epics immunology station following manufacturer's instructions.

Prepara
Hepar
aliquot
lymph
tubes.
and g
was at
pidiun
stainir
Flow
Cells
Coul
throu
electr
cell c
to pr

Prepa
Propag
To ce
suppl
T ly
daily
Harve
Whe
suspe
cent
cell
time
Prepa
Each
soni
mixt
for 1
Esta
The
nor
esta
lym

Preparation of cell cycles

Heparinized whole blood or lymphocytes were used and 100 μ l aliquots of heparinized whole blood or 100 μ l of washed isolated lymphocytes (suspended in normal saline) were placed in 13 x 75 mm tubes. To each tube was added 200 μ l of lysing buffer (Coulter) and gently mixed for ~15 s, then 2.0 ml of DNA stain with RNase was added and mixed for 20 s. The mixture was stained with propidium iodide at a concentration of 50 mg per ml, and allowed a staining period of 15 min and then analysed by flow cytometry.

Flow micro-fluorometric analysis

Cells tagged fluorometrically for DNA content were analysed on the Coulter Epic C flow cytometer, where they were allowed to pass through a laser beam tuned to 488 nm. Fluorescence was measured electronically and recorded as a histogram. DNA distribution in the cell cycle was calculated on accumulated data by parametric analysis to produce a particular DNA histogram.

Preparation of autogenous lymphocytic factor (ALF)

Propagation of lymphocytes

To cell culture flasks, containing 10 ml RPMI 1640 medium and/or supplemented with 1 ml bovine calf serum was added 1 ml of washed T lymphocytes. The culture was incubated at 37°C and monitored daily for approximately 2 to 3 weeks.

Harvesting of cells

When the cell population approximated 5 to 9 x 10⁶ per ml, culture suspensions were transferred to calibrated 50 ml conical tubes and centrifuged for 30 min at 500 x g. The supernate was discarded, cell pellet resuspended in 10 ml normal saline and washed three times by centrifugation for 30 min each time.

Preparation of ALF

Each cell peller was resuspended in ~2.0 ml normal saline and sonicated at 20 W with a duty cycle of 50% for 60 s. The sonicated mixture was sterilized by filtration and stored at -20°C to be diluted for therapeutic use as indicated.

Establishment of a functional T lymphocyte cell cycle

The profile of an individual's T lymphocyte cell cycle may appear normal, yet the lymphocytes may not be functional. A process was established to evaluate the immunological responsiveness of the lymphocytes of the cell cycle.

Stimulation of peripheral lymphocytes

NúncionTM-plastic plates (24 wells) containing 2 ml of RPMI 1640 medium per well were inoculated with ~ 2 to 4×10^6 lymphocytes per ml, 1.25 ng interleukin 1 alpha (Il_1 alpha) per ml, or with 0.1 ml ALF per ml; negative controls had neither Il_1 nor ALF. The plates were incubated at 37°C for 96 h. Cells were harvested, washed and analysed by flow cytometrically as mentioned above. Comparison of the S and G_2 phases of both the unstimulated and the stimulated T lymphocytes was undertaken and an index computed from the equation:

$$\text{stimulation index} = (\text{stimulated S} + G_2) / (\text{unstimulated S} + G_2)$$

Clinical Investigation

A total of 315 individuals was investigated and of these, 25 were controls. T lymphocyte cell cycle profiles, haematological, T and B profiles, and CMI tests were carried out on all candidates. Individuals with immune deregulation were treated with autogenous lymphocytic factor (ALF). These individuals were chemically sensitive and chronically ill. The illnesses include dermatitis, vasculitis, asthma, organic brain syndrome, and Gulf war syndrome. In all cases, these illnesses were characterized with immune system suppression, dysfunction or deregulation.

Major symptoms presented by these patients include one or more of the following symptoms: lacrimation, pruritis, swelling and puffiness (ocular); fullness, noise, and dizziness (otic); congestion, sneezing, rhinorrhoea and blowing (nasal); lump, clearing and postnasal drip (throat); hypersensitivity reactions (immune); arthritis and arthralgia, fatigue and muscle pain (musculoskeletal); pressure and cough (chest); weight loss and fatigue (constitutional); miliary, ethmoidal and frontal (headache); insomnia, shortness of breath and depression (neurological).

All 290 chemically sensitive individuals presented a history of being affected by environmental incitants found in categories such as food, biological inhalants, and chemicals. They presented histories of varied backgrounds, but common among them was that all showed irregular cell cycles and abnormal T lymphocyte profiles in both numbers and functions in T and B lymphocytes and subsets. DNA histograms showed over or under accumulation of various subtypes of lymphocytes in one or more phases of the cell cycle of each individual.

Cell-mediated Immunity

Delayed cutaneous hypersensitivity, or cell-mediated immunity (CMI) responses were evaluated in 190 patients; the results were recorded before and after treatment with ALF. The multitest CMI test kit (Mérieux Institute, Miami, Florida) was used. Each kit contained seven antigens: tetanus, diphtheria, candida, proteus, streptococcus, trichophyton, and tuberculin. The tests were evaluated and read at 48 h. Evaluation involves scoring the size, and number of increase of the wheals. The diameter of each induration was measured in millimetres and averaged. A reaction was considered positive if the average diameter was 2 mm or more.

Comparative evaluation of T and B lymphocytes

T and B lymphocytes and subsets were evaluated flow cytometrically before and after treatment with ALF.

Table 1 Representative profile of symptoms and signs after autogenous lymphocytic factor treatment

Symptoms	Improvement:		No change:		Deterioration:		Total patients
	Number	%	Number	%	Number	%	
Hypersensitive reaction	63	63	33	33	1	1	100
Recurrent Infection	38	57	29	43	0	0	67
Fatigue	60	68	27	31	1	1	88
Lack of concentration	43	54	36	46	0	0	79
Arthritis	19	44	23	54	1	2	43
Gastrointestinal upset	29	40	43	60	0	0	72
Headache	28	44	33	53	2	3	64
Depression	42	58	30	42	0	0	72

Establishment of therapeutic dosage of ALF

In vitro and *in vivo* procedures were used to establish dosage. Blastogenesis was the *in vitro* test employed for establishing dosage in general. *In vitro* grown peripheral lymphocytes were challenged comparatively with the known mitogens phytohaemagglutinin (PHA), concanavalin A (Con A), and pokeweed mitogen (PWM) of varied concentrations. A dose responsive curve was established to evaluate the efficacy of ALF with that of the mitogens; 0.1 ml of 1:10 dilution of ALF was found to be optimal when administered twice weekly. Intradermal test was carried out on all recipients depending on their response and sensitivity. If unusual sensitivities were observed, the ALF was further diluted. This procedure was successively repeated until no adverse reaction was observed.

Results

Significant changes were observed within 1 to 6 weeks in patients treated with ALF. Changes were observed in immune regulation and overall clinical manifestations. With regard to clinical manifestations, there were noteworthy improvements (Table 1), although minimal symptoms continued after approximately 3 weeks of continued therapy.

In a normal cell cycle, the highest percentage of lymphocytes should be in the G_0 to G_1 phase (Figure 1). This percentage will change dramatically when these lymphocytes are stimulated by various incitants (Figures 3a and 3b). Consequently, different percentages will appear in the S and G_2 M phases, producing a deregulated profile (Figures 1 to 4). Treatment with ALF regulates the T lymphocyte cell cycle profile.

Immunologically, there were significant regulations of T lymphocyte cell cycles, especially from one phase of the cycle to another. Patients became less sensitive and more tolerant to specific incitants. As treatment continued, in general in about 6 weeks, a more drastic shift toward that of a normal profile was observed. Figures 1 to 4 summarize the regulatory changes of some of the cell cycles studied.

Changes were observed in the profiles of the T and B cells where T and B lymphocytes and their subsets were evaluated before and after treatment. There was a significant ($p < 0.01$) change in the total lymphocyte count and subsets, T_4 , T_8 in 92 patients investigated. It should be noted that ALF seems to act as an immune modifier since the total lymphocytes, T_4 and T_8 were significantly elevated or reduced in order to maintain normalization (Tables 2 and 3).

1 dosage.
dosage in
ged com-
1 (PHA),
of varied
evaluate
) dilution
e weekly.
on their
ved, the
repeated

patients
gulation
l mani-
lthough
eks of

hocytes
age will
various
ges will
profile
yte cell

mpho-
nother.
sitants.
drastic
l to 4
died.
where
e and
e total
gated.
odifier
vated
id 3).

et al.




Figure 1 Diagrammatic representation of a normal mammalian cell cycle, showing the relative number of lymphocytes in each phase of the cycle, as displayed by the intensity of fluorescence of the fluorochrome-bound DNA in each lymphocyte. Normally, the percentage of lymphocytes in each phase of the cell cycle is 90-95, 5-10 and 0-5 in the G₀G₁ phase, S phase and G₂M phase, respectively. These quiescent lymphocytes display, normally, a low level of replication, and progression through the cell cycle. In response to stimuli, these cells undergo rapid division and cycle rapidly. The shadings show that there is overlapping of phases and not specific lines of demarcation: stippled, G₀G₁ phase (90-95% noncycling lymphocytes); hatched, S phase (5-10% cycling lymphocytes); unshaded (open), G₂M phase (0-5% cycling lymphocytes); and solid areas indicate overlapping of phases.

Table 2 Representative changes in T and B cell profiles after 92 chemically sensitive patients were treated with autogenous lymphocytic factor

Treatment	Total lymphocytes	T ₁₁	T ₄	T ₈	T ₄ /T ₈	B ₄
Before	2,112 ± 632	1,624 ± 457	930 ± 45	439 ± 58	2.3 ± 0.8	188 ± 102
After	2,232 ± 676*	1,634 ± 544	1,027 ± 297**	478 ± 189**	1.3 ± 7.4	171 ± 120

* p <0.05; **p <0.01; n = 92.




Figure 2 A representative DNA histogram of peripheral T lymphocytes from normal volunteers. Note that the highest percentage of lymphocytes is in the G_0G_1 phase, as expected, but the percentage varies in different individuals in keeping with the prevalence of environmental pollutants. Thus, the ideal normal profile is seldom achieved. The abscissa shows the relative fluorescence of the fluorochrome-bound DNA in each cell.

Table 3 Profile of T lymphocyte subsets modulation after treatment with autogenous lymphocytic factor

After treatment	Total lymphocytes:		T_{11}		T_4		T_8	
	Number	%	Number	%	Number	%	Number	%
Increase	52	57	41	46	53	53	55	60
Decrease	40	43	51	54	39	42	37	40
Probability	<0.05		>0.05		<0.05		<0.01	

Table 4a Typical cell-mediated response in (number and size) chemically sensitive individuals treated with autogenous lymphocytic factor

Patients	Before ALF	After ALF	P
180	6.46 ± 5.81	9.28 ± 7.25	<0.001

Results are given as mean ± SD.

Patients showed significant improvement ($p < 0.001$) in their CMI scores (Table 4). The regulatory effect of the immune system can be objectively assessed periodically after the initial treatment with ALF. The resourceful parameters are profiles of T lymphocyte cell cycle, T and B lymphocytes, and their subsets, cell-mediated immunity, signs and symptoms.

Side effects of ALF

The side effects were minimal and occurred only in six adults (five females and one male) where the average age was 52 years. These patients were intolerant of ALF, their symptoms included pain and irritation in the throat, burning in the eyes, pain and irritation in the chest, heat palpitations, influenza-like symptoms, headache, fatigue, and chills.

Table 4b Cell-mediated immunity positive score (number and size) after treatment with autogenous lymphocytic factor

Score	Patients	%	P
Increase	58	74	<0.001
No change	9	12	
Decrease	11	14	




Figure 3a A representative irregular T lymphocyte cell cycle profile showing the effect(s) of stimulating the environmentally ill patients influence, primarily, the lymphocytes in the S phase.

Figure 3b A representative irregular T lymphocyte cell cycle profile showing the effect(s) of the stimulating incitants on the lymphocytes in the G₂M phase. It seems that each incitant, or a mixture of incitants affects lymphocytes in a particular phase(s) of the cell cycle, resulting in a variety of irregular cell cycle profiles and presumably dictates varied patterns of clinical manifestations. The abscissae in Figures 3a and 3b show the relative fluorescence of the fluorochrome-bound DNA in each cell.

Discussion

The flow cytometric profile of the T lymphocyte cell cycle as demonstrated by DNA histogram presents a reflection of the status of T lymphocytes in an individual. Of great importance is that its application is not limited to a certain category of individuals, but to normal subjects as well as individuals who are compromised by varied incitants. The essence of its clinico-biological importance, as detected in the present investigation, is that it is a reflection of T lymphocyte cell function, and may facilitate an in-depth approach to the treatment of some immunodeficient illnesses.

The progression of T lymphocytes from one phase of the cell cycle to another is time-dependent, ~8 to 12 h in the G₀-G₁ phase, 6 to 8 h in the S phase, and 0.5 to 1 h in the G₂M phase. The 290 individuals who were investigated in this study were affected principally by environmental incitants in categories such as foods, biological inhalants, and chemicals. They presented histories of varied backgrounds and different cell cycle profiles. The DNA histogram




Figure 4a Initial T lymphocyte cell cycle profile of a patient on examination at the clinic. Note that the irregularity of the profile is emphasized in the G₂M phase.

Figure 4b Cell cycle of the patient in Figure 4a after the subject was treated with autogenous lymphocytic factor (ALF) for 3 weeks. Note that the percentages of lymphocytes in each phase of the cell cycle are still irregular, but there has been a significant improvement. The highest percentage of lymphocytes now appears in the G₀G₁ phase.

Figure 4c T lymphocyte cell cycle of patient (Figures 4a and b) subsequent to 3 months treatment with ALF. Note that the cell cycle has attained a relatively normal profile. ALF seems to act as a biological response modifier. The abscissae in Figures 4a, b and c show the relative fluorescence of the fluorochrome-bound DNA in each cell.

cycles showed over or under accumulation of lymphocytes in one or more phases of the cell cycle of each individual (Figures 3a and 3b). As shown in Figure 2, even the normal controls do not present an ideal DNA histogram profile of T lymphocytes. This is due to the fact that, in general, the ideal environment is seldom ever achieved, thus there is always some degree of immunological compromise to most individuals. It is reasonable then to assume that the incitant(s) which lead to immuno-incompetence of these individuals are capable of inactivating the particular enzyme(s), cyclin-dependent kinases, or depleting specific cyclins whose combination with their specific kinases are instrumental in catalysing the progression of T lymphocytes from one phase to another of the cell cycle. The T lymphocytes of the individual affected are locked in a particular phase, either resting, synthesizing, or multiplying too much in the G_0 - G_1 , S, G_2 M phase respectively. Thus the subject manifests symptoms peculiar to the phases(s) affected. This hypothesis offers an opportunity to associate clinical manifestations with T lymphocyte cell cycle irregularity.

The treatment of lasting importance would be reasonably thought of as a biological response modifier, which would stimulate CDKs, regulate the cell cycle and the enzymes of purine and pyrimidine nucleotide synthesis. It is now generally accepted that these enzymes are elevated during the S phase (Cory, 1993). It seems logical that ALF stimulates these regulatory functions. The autogenous lymphocytic factor is a dialysable mixture of the many effector substances which may be released from *in vitro* grown stimulated lymphocytes with the ability to invoke immunological influences *in vivo* or *in vitro*, (Griffiths and Rea, 1995). The expected biological activity(s) of ALF would be to act as a biological response modifier, with mechanisms of action as suppressor and regulator of the immune system, especially in the regulation of the T lymphocyte cell cycle.

Kamp *et al.* (1994) showed that progression through the cell cycle requires the joint influence of positive regulator subunits, cyclins, and cyclin dependent kinases. These subunits regulate by phosphorylation a number of key substrates which subsequently activate a transition from G_1 to S, and G_2 to M phases of the cell cycle. Phosphate groups are transferred from ATP to a special amino acid in the target protein by protein kinases, while phosphatases remove the phosphate groups from the target proteins. The addition and removal of phosphate groups significantly affect the biochemical behaviour of the target proteins. Many protein kinases and phosphatases have a specific affinity for their target proteins, and act as determinants for controlling the activity(s) of their target

proteins. Indeed, ALF may possess these protein kinases which, acting as molecular switches, regulate an irregular T lymphocyte cell cycle to that of an ordered and orderly progression.

As demonstrated (Figure 4c), ALF regulates deregulated cell cycles, improves immunological profiles by increasing or decreasing the number of circulating lymphocytes and their subpopulations (Tables 2 and 3); and restores immune responses as demonstrated by enhanced cutaneous hypersensitivity (CMI) (Table 4). ALF also invoked immediate intradermal test response within 45 min of administration subsequent to the administration of antigen challenges which were previously negative or delayed for at least 15 days. Responses were observed by the number and size of wheals to the antigens which were inoculated. It is noteworthy that some chemically sensitive patients responded favourably, or completely recovered after environmental exposure to air, food, water, nutritional support and exercise. Some patients responded but did not fully recover. The patients who recovered reacted profoundly to exposure of very minute environmental insults of ambient chemicals, biological inhalants, and foods, or inhalants. However, when these patients were treated with ALF, their hypersensitivity was markedly reduced or disappeared; there was also significant improvement in recurrent infection, fatigue, headaches, depression, concentration, and gastrointestinal upsets. Eight patients immediately improved on initial injection with ALF.

T-suppressor, T-helper cells, and total lymphocytes were increased as reflected in an increase in a low population, and a decrease in a too high population. This suggests that ALF is a modulator rather than a stimulator. The mechanism(s) of action of ALF have not yet been ascertained. Indeed, the purification and determination of ALF from sorted T lymphocytes with or without helper phenotypes will be necessary to facilitate understanding the mechanism of action. However, it has been documented that the human cells contain a regulatory protein, CKS protein, which is a genetic suppressor of temperature sensitive CDK mutations. There are two isoforms of this CKS protein, namely CKSH1 and CKSH2. It is believed that CKSH2 protein binds to the catalytic subunit of the CDKs, and is essential for their biological function (Parge *et al.*, 1993). To this end, further investigations are being carried out.

The concept of a functional cell cycle is timely. Indeed, the profile of a cell cycle offers an insight into the immune status of a patient. However, the cell cycle profile does not indicate whether or not the lymphocytes are immunologically responsive or capable of responding to therapy. The influence of varied environmental incitants and/or

other systemic illnesses may render these lymphocytes incapable of normal progression through the cell cycle, thus resulting in a non-functional cell cycle. *In vitro* lymphocyte activation represents a standard procedure for evaluating cell mediated responses to a variety of stimuli including antibodies, polyclonal mitogens, specific antigens and cytokines. Interleukin-1 alpha (IL₁ alpha) and ALF were used as activators in the present investigation. Interleukin-1 alpha was used due to its extremely broad spectrum of bioactivity. ALF was used to observe its capability as a biological response modifier. The stimulatory process presented in this investigation offers a method of computing lymphocyte stimulatory response, and consequently an opportunity to establish an individual regime of treatment of illnesses associated with immunodeficiency. This investigation offers: (1) a vivid picture of the status of the immune profile of a patient as demonstrated by the DNA histogram of the T lymphocyte cell cycle; (2) an opportunity to regulate an irregular T lymphocyte cell cycle by treatment with autogenous lymphocytic factor; and (3) a clinical tool for the regulation of abnormal cell cycles and/or haematological and immunological profiles in humans with immunological deregulation. It should be noted that there is a pending United States patent, number 08/380,063 for the application of the cell cycle to clinical medicine, and also the regulation of the cycle by autogenous lymphocytic factor.

Acknowledgements

The authors are grateful to Dr Alfred Johnson and Dr Gerald Ross of the Environmental Health Center - Dallas, for their contribution to the clinical data, and to Dr Vernon E. Scholes for reviewing this manuscript.

References

- CORY J. G. 1993. Purine and pyrimidine nucleotide metabolism. *In* Biochemistry with Clinical Correlation. pp 629-73. Edited by T. M. Devlin. Wiley-Liss, New York.
- CROSS F. R. 1988. DAF 1 a mutant affecting size control, pheromone arrest, and cell kinetics of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 8 4875-84.
- DESAI D., Gu Y. and Morgan D. O. 1992. Activation of human cyclin-dependent kinases *in vitro*. *Mol. Cell. Biol.* 3 571-82.
- DRAETCA G. and Beach D. 1988. Activation of CDC2 protein kinase during mitosis in human cells: cell cycle-dependent phosphorylation and subunit rearrangement. *Cell* 3 17-26.
- EVANS T. E., Rosenthal E. T., Youngblom J., Distel D. and Hunt T. 1983. Cyclin: a protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell* 33 389-96.
- FANG F. and Newport J. W. 1991. Evidence that the G₁-S and G₂-M transitions are controlled by different CDC2 proteins in higher eukaryotes. *Cell* 66 731-42.
- GIVAN A. L. 1992. Cell from within. *In* Flow Cytometry - First Principles. pp 103-33. Edited by A. L. Givan. Wiley-Liss, New York.

GLO1
pathv
GOL1
prote
GRIF1
GUT1
is a s
HAD1
of cy
6255
HEL1
HUN
to a
KAM
Stoc
pote
KINC
557-
LEE
Hen
E. F
LIV1
LEH
Cell
LEW
61
LOR
yeas
ME1
cycl
ME1
199
MU
cell
MU
of t
NA1
111
NU
G₂1
NU
503
NU
PA1
req
PA1
ato
387
PI1
M-
58
RE
pai
RE
ch
C11
RE
Ge

- GLOTZER M., Murray W. A. and Kirschner M. W. 1991. Cyclin is degraded by the ubiquitin pathway. *Nature* 349 132-8.
- GOLD K. L. and Nurse P. 1989. Tyrosine phosphorylation of the fission yeast CDC2+ protein kinase regulated entry into mitosis. *Nature* 324 39-45.
- GRIFFITHS B. B. and Rea W. J. 1995. Pending U.S. patent number 08/380,063.
- GUTIER J., Solomon M. J., Booher R. N., Bazan J. F. and Kirschner M. W. 1991. CDC25 is a specific tyrosine phosphatase that directly activates p34 CDC2. *Cell* 67 197-211.
- HADWIGER J. A., Wittenberg C., Richardson H. E., Lopes M. and Reed S. I. 1989. A family of cyclin homologs that control the G phase in yeast. *Proc. natn. Acad. Sci. U.S.A.* 86 6255-8.
- HELCHMAN K. A. and Roberts J. M. 1994. Rules to replicate by. *Cell* 79 557-62.
- HUNTER T. and Pines J. 1994. Cyclins and cancer 11: cyclin D and CDK inhibitors come to age. *Cell* 79 573-82.
- KAMP A., Grula N. A., Weaver-Feldhaus J., Liu Q., Harshman K., Tavtigian S. V., Stockert E., Day R. S. III, Johnson B. E. and Skolnick M. J. 1994. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 264 436-40.
- KING R. W., Jackson P. K. and Kirschner M. W. 1994. Mitosis in transition. *Cell* 79 557-62.
- LEE W. M. and Dang C. V. 1995. Control of cell growth, differentiation and death. In *Hematology, Basic Principles and Practice*. pp 89-84. Edited by R. Hoffman, E. KaBenz Jr, S. J. Shattil, B. Furie, H. J. Cohen and L. E. Silberstein. Churchill Livingstone, New York.
- LEHNER C. and O'Farrell P. 1990. The rule of *Drosophila* cyclin A and B in mitotic control. *Cell* 61 535-47.
- LEWIN B. 1990. Driving the cell cycle: M phase kinase, its partners, and substrates. *Cell* 61 743-52.
- LORINEZ A. T. and Reed S. I. 1984. Primary structure homology between the product of yeast cell division control gene CDC-28 and vertebrate oncogenes. *Nature* 307 183-5.
- MEYERSON M. and Harlow E. 1994. Identification of G₁ kinase activity for CDK6, a novel cyclin D partner. *Mol. Cell. Biol.* 14 2077-86.
- MEYERSON M., Enders G. H., Wie C. L., Su L., Gorka C., Nelson C., Harlow E. and Tsai L. 1992. A family of human CDC2-related protein kinases. *EMBO J.* 11 2909-17.
- MURRAY A. W. and Kirschner M. W. 1989. Cyclin synthesis drives the early embryonic cell cycle. *Nature* 339 275-80.
- MURRAY A. W. and Kirschner M. W. 1989. Dominoes and clocks: the union of two views of the cell cycle. *Science* 248 614-21.
- NASMYTH K. A. 1990. Far-reaching discoveries about the regulation of START. *Cell* 63 1117-20.
- NURSE P. and Bissett Y. 1981. Gene required in G₁ for commitment to cell cycle and in G₂ for control of mitosis in fission yeast. *Nature* 292 558-60.
- NURSE P. 1990. Universal control mechanism regulating onset of M-phase. *Nature* 344 503-8.
- NURSE P. 1994. Ordering S phase and M phase in the cell cycle. *Cell* 79 547-50.
- PAGANO M., Pepperkok P., Verde F., Ansorge W. and Draetta G. 1992. Cyclin A is required at two points in the human cell cycle. *EMBO J.* 11 961-71.
- PARGE H. E., Arvai A. S., Murtari D. T., Reed S. I. and Telner J. A. 1993. Human CksHs2 atomic structure: a role for its hexameric assembly in cell cycle control. *Science* 262 387-95.
- PINES J. and Hunter T. 1989. Isolation of human cyclin C DNA: evidence for cyclin M-RNA and protein regulation in the cell cycle and for interaction with P34 CDC2. *Cell* 58 833-46.
- REA W. J., Pan Y., Johnson A. R., Fenyves E. J. and Smadi N. 1986. T and B lymphocyte parameters measured in chemically sensitive patients and controls. *Clin. Ecol.* 4 11-4.
- REA W. J., Pan Y., Johnson A. R. and Fenyves E. J. 1987. T and B lymphocytes in chemically sensitive patients with toxic volatile organic hydrocarbons in their blood. *Clin. Ecol.* 5 171-5.
- REED S. I. 1992. G₁ specific cyclins in search of an S-phase-promoting factor. *Trends Genet.* 7 95-9.

- RICHARDSON H. E., Wittenberg C., Cross F. and Reed S. I. 1989. An essential G₁ function for cyclin-like proteins in yeast. *Cell* 59 1127-33.
- SHERR C. J. 1993. Mammalian G₁ cyclins. *Cell* 73 1059-65.
- SHERR C. J. 1994. G₁ phase progression: cycling on cue. *Cell* 79 551-5.
- SOLOMON M. J., Glotzer M., Lee T. H., Philippe M. and Kirschner M. W. 1990. Cyclin activation of p34 CDC2. *Cell* 63 1013-24.
- WALKER D. H. and Maller J. L. 1991. Role of cyclin A in the dependence of mitosis on completion of DNA replication. *Nature* 354 314-7.
- WANG J., Chenevessé X., Hanglein B. and Brechot C. 1990. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 343 555-7.
- WITTENBERG C., Sugimoto K. and Reed S. I. 1990. G₁-specific cyclins of *S. cerevisiae*: cell cycle periodicity, regulation by mating pheromone, and association with the p34 CDC28 protein kinase. *Cell* 62 225-37.
- XIONG Y., Connolly T., Fletcher B. and Beach D. 1991. Human D-type cyclin. *Cell* 65 691-9.
- ZINDY F., Lamas E., Chenevessé X., Sobczak J., Wang J., Fasquet D., Hanglein B. and Brechot C. 1992. Cyclin A is required in S phase in normal epithelial cells. *Biochem. biophys. Res. Commun.* 182 1144-54.

Accepted 20 January 1993



241057

STIC-ILL

From: Schwadron, Ronald
Sent: Wednesday, February 10, 1999 11:02 AM
To: STIC-ILL
Subject: 08902692

Youdim et al.
Clin. Ecology
vol 7
p 55-61
1990

Youdim et al.
Clin. Ecology
vol 7
p 62-66
1990

Youdim et al.
Clin. Ecology
vol 7
p 67-72
1990

Rea
Annual Meeting-American Academy of Environmental Medicine (I think this is a book of abstracts published by American academy of Environmental Medicine)
Sept 1995
p213-222

0735-9306

Griffiths et al.
Cytobios
vol 93
p49-66
1998

thanxs

ask
2/11

nlm
2/19
604

2/10
ND

NIH ACS
2/10 - RC

TREATMENT OF ENVIRONMENTALLY SENSITIVE PATIENTS WITH TRANSFER FACTOR

PART II: CLINICAL STUDIES AND IMMUNOLOGICAL CORRELATES

Said Youdim, Ph.D.¹
William J. Rea, M.D.²

ABSTRACT

Fifty patients ranging in age from 7 to 75 with hypersensitivity reactions to foods, inhalants and chemicals having symptoms of cephalgia, recurrent infections, fatigue, lack of concentration, arthritis, gastrointestinal problems and depression were treated with a course of transfer factor one unit two times per week for a period of 6-12 months. Seventy-eight percent of the patients had some relief as indicated by decrease in both the severity and frequency of one or more of their symptoms. These clinical improvements correlated well with increased cutaneous delayed type hypersensitivity (cellular immune response) (73.5%), as well as increase in the total number of lymphocytes (66.7%), total T cells (66.7%) and T helper cells (70.4%). These data indicate that antigen specific and possibly non-specific components of TF may modify the immunologic response of TF recipients and improve their clinical status. *Transfer factor, cellular immune response, T & B Lymphocytes.*

¹11927 Darlington Ave, #4
Los Angeles, CA 90049

²Environmental Health Center - Dallas

INTRODUCTION

The clinical use of transfer factor (TF) in the therapy of a wide variety of disorders is well documented (1,2). These include congenital immunodeficiencies, acquired immunodeficiencies, malignancies, autoimmune and neurological, as well as infectious diseases such as protozoal, mycobacterial, fungal and viral diseases. In most cases the effect of TF on such infections is prophylactic and may result in either the initiation or restoration of cellular immunity. Lawrence reported that sixty-six percent (24/36) of patients with congenital immunodeficiency given TF converted to positive cutaneous delayed type hypersensitivity (DTH), and/or produced migration inhibitory factor (MIF), and/or responded to the corresponding antigens by lymphocyte transformation. In addition, 44% (16/36) of his patients improved clinically (1). In these studies, the transfer of an antigen-specific moiety, as well as nonspecific components in the TF preparation, was demonstrated. In fact, some studies (3,4) have shown that at least 130 different moieties are present in the extract prepared by the method of Lawrence. Fudenberg, et. al. (5) described some of these as a nonspecific adjuvants of cell mediated immunity while other components such as hypoxanthine, may be inhibitory.

We treated 50 environmentally sensitive patients, some with immune dysregulation, with a preparation of TF of molecular weight up to 30,000 Dalton units. Since the antigen-specific component of TF is reported to have a molecular weight between 3,500 and 12,000 Dalton units (6), it can be safely assumed that the above preparation also contained nonspecific components.

MATERIAL AND METHODS

Preparation of transfer factor (TF), lymphocyte phenotyping, and performance of cell mediated immunity (CMI) tests were as described in the accompanying paper (7).

Patient Selection: Patients with food allergies (8,9),

chemical, and environmental sensitivities (10) were advised to adopt environmentally safe habits such as living in a natural gas-free environment and to use less chemically contaminated food and water. The majority of patients also took antigen immunotherapy for foods, inhalants, or chemicals (8,11,12) concurrently with TF.

Patient Questionnaire: Each patient was given a "Symptoms Score Sheet" to be filled in prior to and following 6-12 months of TF therapy. The patients were asked to respond as to the frequency and severity of symptoms in the following categories: hypersensitivity reactions to incitants, cephalgia, recurrent infections, fatigue, gastrointestinal problems, depression, and lack of concentration. Based on the patient's response on a scale of 1 to 5, each respondent was categorized as demonstrating moderate decrease, substantial decrease, (i.e. improvement), or no change in his/her symptoms. The graded scores of frequency and severity of symptoms were evaluated before and after therapy.

TF Dose: Each patient received two weekly doses of TF each, injected subcutaneously or intramuscularly for 6-12 months. Each dose of 1 ml contained 10^8 lymphocyte equivalents.

RESULTS

Fifty patients were entered into a program of TF therapy for 6 to 12 months. Table 1 shows the distribution of age and sex among these patients. There were a total of 8 males and 42 females ranging in age from 7 to 75 years. Each patient was asked to submit a symptoms score before and after TF therapy. Tables 2 and 3 summarize data generated from these questionnaires. Depending on each symptom category, patients showed either a moderate decrease, substantial decrease, or no change in the severity or frequency of their symptoms. A substantial number of patients admitted to some relief as indicated by decrease in both severity and frequency of their symptoms.

We next examined the correlation of improved clinical status to the increase in the number of WBC, lymphocytes, T cells, T cell subpopulations, B cells and CMI response in TF recipients. These data are given in Tables 4 and 5. 39 of 50 patients (78%) responded to TF therapy with some improvement in one or more of their symptoms (Tables 2, 3 and 5). The closest immunologic correlate of improvement in clinical status was the patient's CMI response (73.5%), followed by increase in the number of T helper cells (70.4%). Both improvement in CMI and T helper cell numbers show excellent correlation to the decrease in the frequency of recurrent infection of 70.6% (Table 3). In every case, an increase in the number of WBC, total number of lymphocytes, T11, T4, T8, and B

lymphocytes were in good agreement with improvement in clinical status.

Not all patients, however, responded well to TF therapy. In fact, 22% of the patients demonstrated no improvement in their clinical status and a minor population showed no change or a decrease in their immune parameters (Table 4).

The cumulative data indicates that a substantial portion of the patients (73.5%) exhibited an increase in their CMI, in the number of their B and T lymphocytes (62.1%) or in both of these parameters (95.5%) (Table 4).

TABLE 1
The Distribution of Age and Sex
Among 50 Patients Treated with TF

AGE/SEX	MALE	FEMALE	TOTAL
01-10	1	00	01
11-20	1	01	02
21-30	2	02	04
31-40	1	14	15
41-50	3	10	13
51-60	0	11	11
61-70	0	03	03
71-80	0	01	01
TOTAL	8	42	50

DISCUSSION

The results presented in this paper (Tables 2 and 3) indicate improvements in eight clinical parameters reported by the patients enrolled in a program of TF therapy for 6 - 12 months.

A large number of the patients concurrently utilized antigen immunotherapy (8,11,12) and had adopted environmentally less-toxic or less-allergenic living habits. It can also be seen that these same patients were, for the most part, middle-aged females who were long-term referrals at the Environmental Health Center - Dallas (EHC-Dallas).

Improvement of 60% to 70% in various symptom categories by these patients closely paralleled enhanced immunologic parameters (Tables 4 and 5). These results clearly resemble previously published clinical improvement of 44% and in vitro and in vivo cellular immune responses by 66% of congenitally immunodeficient patients treated with TF (1). An interesting question posed at this juncture is the relative contribution of the specific and nonspecific components of the TF preparation. The chemical complexity of the TF preparation (with over one hundred different molecules) makes it difficult to assign specific functional activity to each molecule. Resolution of this

TABLE 2
Improvement in Clinical Severity of Symptoms Indicated by
Patients Receiving Transfer Factor Therapy

	SEVERITY						Total # Patients Improved	Total # Patients Tested	Patients Improved %
	Moderate Decrease		Substantial Decrease		No Change				
	#	%	#	%	#	%			
Hypersensitivity Reactions	27	54	6	12	17	34	33	50	66
Cephalgia	13	31.7	9	21.95	19	46.3	22	41	53.7
Recurrent Infection	16	36.4	10	22.7	18	40.9	26	44	59.1
Fatigue	19	38.8	15	30.6	15	30.6	34	49	69.4
Lack of Concentration	16	37.2	12	27.9	15	34.9	28	43	65.1
Arthritis	9	33.3	3	11.1	15	55.6	12	27	44.4
G.I. Problems	12	41.4	3	10.3	14	48.3	15	29	51.7
Depression	12	37.5	7	21.9	13	40.6	19	32	59.4

TABLE 3
Improvement in Frequency of Symptoms Indicated by
Patients Receiving Transfer Factor Therapy

SYMPTOMS	FREQUENCY						Total # Patients Improved	Total # Patients Tested	Patients Improved %
	Moderate Decrease		Substantial Decrease		No Change				
	#	%	#	%	#	%			
Hypersensitivity Reactions	16	35.6	10	22.2	19	42.2	26	45	57.8
Cephalgia	07	20.6	12	35.3	15	44.1	19	34	55.9
Recurrent Infection	14	41.2	10	29.4	10	29.4	24	34	70.6
Fatigue	14	31.8	13	29.5	17	38.6	27	44	61.3
Lack of Concentration	16	41	11	28.2	12	30.8	27	39	69.2
Arthritis	08	30.8	6	10.7	17	60.7	14	26	53.8
G.I. Problems	08	28.6	03	10.7	17	60.7	11	28	39.3
Depression	10	32.3	08	25.8	13	41.9	18	31	58.1

TABLE 4
Changes in Immune Parameters After Transfer Factor Therapy

Immune Parameters	# Patients Tested	Below Normal Range Change to Normal Ranges		Partial Increase Towards Normal Range		Total Increase in Range		No Change		Decrease	
		#	%	#	%	#	%	#	%	#	%
CMI*	34	12	35.3	13	38.2	25	73.5	05	14.7	04	11.8
T&B**	29	09	31.03	09	31.03	18	62.1	09	31.03	02	06.9
CMI & T&B	22	09	40.9	12	54.5	21	95.5	01	04.5	0	0
CMI in Normal Population		Mean # Reactions				Mean Reaction Size					
315 Male		4.5				18.3					
299 Female		3.5				12.2					

*Data provided by Institute Merieux, Lyon, France

**Normal ranges established on 60 males and females by EHC-Dallas

difficulty will have to await the isolation and purification of each indigenous molecule. Such components as thymosin A-1 like materials (13), Interferon (14), prostaglandins (15), serotonin, ascorbate, chemoattractants and immobilizing factors (1,5) must exert considerable immunological, biochemical, hormonal and other physiological stimuli on the TF recipient. It is well known, for example, that interferon (16) exhibits at least some 15 enzymatic and biochemical properties and 27 immunoregulatory functions. The anti-viral (17,18) and antiproliferative (19) effects of interferon are well known and could be responsible for decreased recurrent infections. Thymosin A-1 like materials (13) enhance Migration Inhibitory Factor (MIF), Alpha and Gamma interferon, Interleukin-2, and Lymphotoxin production, and stimulate viral, fungal and tumor immunity and increases IL-2 receptor expression as well as amplify T-cell immunity in humans (20,21).

All of these effects modulate biological and immunoregulatory functions, and may be responsible for the changes observed in our patients. Prostaglandins (PG), another component of TF (15), are a family of biologically active compounds (22) and products of arachidonic acid metabolism via a cyclo-oxygenase dependent pathway. These compounds are vital components of the balance achieved by normal immunoregulatory mechanisms and implicated in inflammatory reactions, via PG-producing monocytes. PGE2 modulates T-cell activation, T-cell response to antigen, lymphokine production, B-cell and natural killer cell activity (23). Serotonin and histamine are bioactive amines (24,25) and components of the immediate or type I allergic reaction.

TABLE 5
Correlation of Increase in the Number of Lymphocytes and CMI Response Versus Clinical Status After Transfer Factor Therapy

	# Patients Tested	# Patients Improved	%
WBC	27	18	66.7
LYMPHOCYTES	27	18	66.7
T11	27	18	66.7
T4	27	19	70.4
T8	27	15	55.6
T4/T8	27	07	25.9
B LYMPHOCYTES	24	14	58.3
CMI	34	25	73.53
CLINICAL STATUS*	50	39	78.0

*Based on Patient response to:

Hypersensitivity Reactions	Arthritis	Cephalgia
Lack of Concentration	GI Problems	Depression
Recurrent infection	Infection	Fatigue

It is, therefore, possible that both specific (1) and nonspecific modulators (26,27) or "nonimmunologic" components of TF contribute to the improvements observed in the recipients of TF. Although an interesting question, the contribution of each individual moiety will have to await further research. Suffice to say that pooled normal TF, comprised of random subsets of specificities, is a valuable therapeutic adjunct for treatment of environmentally ill patients.

Finally, we should mention that TF was not universally tolerated. A small number of patients were taken off therapy for nonspecific complaints, as well as muscle and joint aches, "cold or flu-like" symptoms, slight fever and local edema or pain at the site of injection. Most of these effects may be attributed to interferon (16) and were transitory in nature.

Alternatively, immunogenetic factors, extreme sensitivity of the patients, quality of the TF preparation or psychologic factors may have played a role in patient intolerance. For the majority of the patients, however, TF was without complications and quite safe. In a double-blind study (Fudenberg and Segre, private communications), showed that long-term administration of TF to mice from time corresponding to middle age until time of death, had no adverse effect upon life span, in fact, mean, median, and maximum life span was the same for the mice that received TF as for those that received saline. Ottgen, et.al.(1, 28) injected pooled TF into women with breast cancer. Doses as large as 257 ml (equivalent to 217 billion cells) were given repeatedly (125 injections) over a period of 310 days. This immunotherapeutic approach testifies to the margin of safety that is afforded by transfer factor.

REFERENCES

1. Lawrence HS. Transfer Factor in Cellular Immunity. *Harvey Lect* 1974;68:239-343.
2. Fudenberg HH, Wilson GB, Goust JM, et al. Dialyzable Leukocyte Extracts (Transfer Factor): A review of clinical results and immunological methods for donor selection, evaluation of activities, and patient monitoring. In: Aituti F, Wigzell H (Eds). *Thymus, Thymic Hormones, and T lymphocytes*. London, Academic Press, 1980:391-421.
3. Wilson GB, Paddock GV, and Fudenberg HH. The Chemical Nature of the Antigen - Specific Moiety of Transfer Factor. *Trans Assoc Am Phys* 1979;92:239-256.
4. Fudenberg HH, Streekauskas AJ, Goust JM, et. al. "Discoid" Lupus Erythematosus: Dramatic Clinical and Immunological Response to Dialyzable Leukocyte Extract (Transfer Factor). *Trans Assoc Am Phys* 1981;94:279-291.
5. Fudenberg HH. Clinical response to "Transfer Factor: Therapy" an update. *Clin Immun Newsletter* 1984;5:109-113.
6. Borkowsky W and Lawrence HS. Deletion of Antigen - Specific Activity from Leukocyte Dialysates Containing Transfer Factor by Antigen - Coated Polystyrene. *J Immunol* 1981;126:486-489.

7. Youdim S, Rea WJ, and Liang HC. Treatment of Environmentally Sensitive Patients with Transfer Factor. I. Immunological Studies *Clin Ecol* 1990;7:55-61.
8. Miller JB. *Food Allergy, Provocative Testing, and Injection Therapy*. Springfield, Thomas, 1972.
9. Brostoff J, Challacombe SJ, (Eds). *Food Allergy and Intolerance*. London, Bailliere Tindall, 1987.
10. Rea WJ, Mitchell BA. Chemical Sensitivity and the Environment. *Immunol and Allergy Practice* 1982;4:5:21-31.
11. King PK, Rubin WA, Fadal RG, et al. Provocation-Neutralization: A Two-Part Study - Part I. The Intracutaneous Provocative Food Test: A multi-Center Comparison Study. *Otolaryngol Head and Neck Surgery* 1988;99:263-271.
12. King PK, Fadal RG, Ward WA, et al. Provocation-Neutralization: A Two-part Study - Part II. Subcutaneous Neutralization, Neutralization therapy: A multi-center study. *Otolaryngol Head Neck Surg* 1988; 99:272-277.
13. Kirkpatrick CH, Khan AA, McClure AL, et al. Thymosin Alpha-1 Like Materials in Dialysates of Leukocyte Extracts. In: Kirkpatrick CH, Burger DR and Lawrence HS, (Eds). *Immunology of Transfer Factor*. New York, Academic Press, 1983:413-420.
14. Khan A, Boettcher J, Garrison O, et al. In Vitro Induction of antiviral Activity by Immunopeptide and Transfer Factor. In: Rainer H and Schattaur FK, (Eds). *Immunotherapy of Malignant Disease*. Verlag Stuttgart, 1978:364-372.
15. Sandler JA, Smith TK, Manganiello C, et al. Stimulation of Monocyte CGP by Leukocyte Dialysates. *Clin Invest* 1975;56:1271-1279.
16. Warren SL. Interferon Review and Clinical Experience. *Immunol Allergy in Pract* 1986;115:33-128.
17. Stevens DA, Merigan TC. Interferon Antibody and Other Host Factors in Herpes Zoster. *J Clin Invest* 1972;51:1170-1178.
18. Dianorzi F, Baron S. Activation by Interferon of the Events Leading to the Antiviral State. *Texas Rep Biol Med* 1977;35:297-306.
19. Gressor I. On the Varied Biological Effects of Interferon. *Cell Immunol* 1977;34:406-415.
20. Oats KK, Szein MB, Goldstein AL. Mechanisms of Action of the Thymosins: Modulation of Lymphokines, Receptors, and T-cell Differentiation Antigens. In: Reif E, Schlesinger M, (Eds). *Cell Surface Antigen Thy-1 Immunology, Neurology, and Therapeutic Applications*. New York, Dekker, 1989:273-288.
21. Szein MB, Goldstein AL. Thymic Hormones - A Clinical Update Springer *Semin Immunopathol* 1986;9:1-18.
22. Ninnemann JL. Prostaglandins: Inflammation and Disease. *Immunol Today* 1984;5:6:173-175.
23. Goodwin JS, Ceuppeus J. Regulation of the Immune Response by Prostaglandins. *J Clin Immunol* 1983;3:295-315.
24. Frick OL. Immediate Hypersensitivity. In: Strites DP, Stobo JD, Fudenberg HH, Wells JV, (Eds). *Basic and Clinical Immunology*. 1985:241-270.
25. Galli SJ. New Approaches for Analysis of Mast Cell maturation, Heterogeneity and Function. *Fed Proc* 1987;46:1906-1914.
26. Gottlieb A, Mazirz GA, Tamaki N, et al. The effect of Dialyzable Products from Human Leukocyte Extracts on Cutaneous Delayed Hypersensitivity Responses. *J Immunol* 1980;124:885- 892.
27. Gottlieb AA, Sutcliffe SB, Maziarz GA, et al. Modification of Intradermal Delayed Hypersensitivity by Components of Leukocyte dialysates In: *Immune Regulators in Transfer Factor*. Khan A, Kirkpatrick CH, Hill NO, (Eds). New York, Academic Press 1979:339.
28. Ottgen HF, Old LJ, Farrow JH, et al. Effect of Dialyzable Transfer Factor in Patients with Breast Cancer. *Proc Natl Acad Sci-US* 1974;71:2319-2323.

7/9/125 (Item 3 from file: 164)
DIALOG(R) File 164: Allied & Alternative Medicine (AMED)
(c) 1998 BLHCIS. All rts. reserv.

00146058 THE BRITISH LIBRARY: 9106863

Treatment of environmentally sensitive patients with transfer factor. Part

I: Immunologic studies

Youdim S; Rea WJ; Liang HC

Clin Ecology, VOLUME: 7, ISSUE: 3-4, PAGE: 55-61

1990

DOCUMENT TYPE: Journal Article LANGUAGE: English RECORD TYPE:
Citation

DESCRIPTORS - Key Word : IMMUNITY; HYPERSENSITIVITY

DESCRIPTORS - Minor Term: IMMUNE SYSTEM; PHYSIOPATHOLOGY

7/9/124 (Item 2 from file: 164)
DIALOG(R)File 164:Allied & Alternative Medicine(AMED)
(c) 1998 BLHCIS. All rts. reserv.

00146059 THE BRITISH LIBRARY: 9106864

Treatment of environmentally sensitive patients with transfer factor. Part
II: Clinical studies and immunological correlates

Youdim S; Rea WJ

Clin Ecology, VOLUME: 7, ISSUE: 3-4, PAGE: 62-6
1990

DOCUMENT TYPE: Journal Article LANGUAGE: English RECORD TYPE:
Citation

DESCRIPTORS - Key Word : HYPERSENSITIVITY; IMMUNITY

DESCRIPTORS - Minor Term: IMMUNE SYSTEM; ENVIRONMENT; PHYSIOPATHOLOGY

(Item 1 from file: 164)
DIALOG(R)File 164:Allied & Alternative Medicine(AMED)
(c) 1998 BLHCIS. All rts. reserv.

00146060 THE BRITISH LIBRARY: 9106865

Treatment of environmentally sensitive patients with transfer factor. Part

III: Case studies on three patients

Youdim S; Liang HC; Rea WJ

Clin Ecology, VOLUME: 7, ISSUE: 3-4, PAGE: 67-72

1990

DOCUMENT TYPE: Journal Article LANGUAGE: English RECORD TYPE:
Citation

DESCRIPTORS - Key Word : HYPERSENSITIVITY; IMMUNITY; ENVIRONMENT

DESCRIPTORS - Minor Term: IMMUNE SYSTEM; PHYSIOPATHOLOGY

01474979 INSIDE CONFERENCE ITEM ID: CN014639864
The Potential Uses for Autogenous Lymphocyte Factors (ALF) in
Environmental Medicine

Rea, W.

CONFERENCE: American Academy of Environmental Medicine: The cutting edge
of environmental medicine-Annual meeting; 30th

ANNUAL MEETING- AMERICAN ACADEMY OF ENVIRONMENTAL MEDICINE, 1995; 30th
P: 213-222

The Academy, 1995

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts, presentations,
papers and programme

CONFERENCE SPONSOR: American Academy of Environmental Medicine

CONFERENCE LOCATION: Tucson, AZ

CONFERENCE DATE: Sep 1995 (19950) (19950)

BRITISH LIBRARY ITEM LOCATION: 1087.378000

34 (Item 34 from file: 5)
DIALOG(R)File 5:BIOSIS PREVIEWS(R)
(c) 1999 BIOSIS. All rts. reserv.

03444499 BIOSIS NO.: 000023017587
CONTROLLED STUDIES ON FOOD SENSITIVITY USING INJECTION THERAPY

AUTHOR: REA W J; WILLIAMS M L; SPRAGUE D E
AUTHOR ADDRESS: DALLAS, TEXAS.

JOURNAL: 37TH ACA (AMERICAN COLLEGE OF ALLERGISTS) SCIENTIFIC CONGRESS,
WASHINGTON, D.C., USA, APRIL 4-8, 1981. ANN ALLERGY 47 (2). 1981. 139.
CODEN: ANAEA
DOCUMENT TYPE: Meeting
RECORD TYPE: Citation

)File 5:BIOSIS PREVIEWS(R)
(c) 1999 BIOSIS. All rts. reserv.

11652669 BIOSIS NO.: 199800434400

The role of the T lymphocytic cell cycle and an autogenous lymphocytic factor in clinical medicine.

AUTHOR: Griffiths Bertie B(a); Rea William J; Griffiths Bradley; Pan
Y

AUTHOR ADDRESS: (a)EHC-D Analytical Lab., Environ. Health Cent.-Dallas,
8345 Walnut Hill Lane, Suite 240, Dallas, T, USA

JOURNAL: Cytobios 93 (372):p49-66 1998

ISSN: 0011-4529

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

(Item 1 from file: 34)
DIALOG(R)File 34:SciSearch(R) Cited Ref Sci
(c) 1999 Inst for Sci Info. All rts. reserv.

07717369 Genuine Article#: 200PB Number of References: 207

Title: Functional somatic syndromes

Author(s): Barsky AJ (REPRINT) ; Borus JF

Corporate Source: BRIGHAM & WOMENS HOSP, DIV PSYCHIAT, 75 FRANCIS
ST/BOSTON//MA/02115 (REPRINT)

Journal: ANNALS OF INTERNAL MEDICINE, 1999, V130, N11 (JUN 1), P
910-921

ISSN: 0003-4819 Publication date: 19990601

Publisher: AMER COLL PHYSICIANS, INDEPENDENCE MALL WEST 6TH AND RACE ST,
PHILADELPHIA, PA 19106-1572

Language: English Document Type: REVIEW

Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences; CC CLIN--Current
Contents, Clinical Medicine;

Journal Subject Category: MEDICINE, GENERAL & INTERNAL

Abstract: The term functional somatic syndrome has been applied to several related syndromes characterized more by symptoms, suffering, and disability than by consistently demonstrable tissue abnormality. These syndromes include multiple **chemical sensitivity**, the sick building syndrome, repetition stress injury, the side effects of silicone breast implants, the Gulf War syndrome, chronic whiplash, the chronic fatigue syndrome, the irritable bowel syndrome, and fibromyalgia. Patients with functional somatic syndromes have explicit and highly elaborated self-diagnoses, and their symptoms are often refractory to reassurance, explanation, and standard treatment of symptoms. They share similar phenomenologies, high rates of co-occurrence, similar epidemiologic characteristics, and higher-than-expected prevalences of psychiatric comorbidity. Although discrete pathophysiologic causes may ultimately be found in some patients with functional somatic syndromes, the suffering of these patients is exacerbated by a self-perpetuating, self-validating cycle in which common, endemic, somatic symptoms are incorrectly attributed to serious abnormality, reinforcing the patient's belief that he or she has a serious disease. Four psychosocial factors propel this cycle of symptom amplification: the belief that one has a serious disease; the expectation that one's condition is likely to worsen; the "sick role," including the effects of litigation and compensation; and the alarming portrayal of the condition as catastrophic and disabling. The climate surrounding functional somatic syndromes includes sensationalized media coverage, profound suspicion of medical expertise and physicians, the mobilization of parties with a vested self-interest in the status of functional somatic syndromes, litigation, and a clinical approach that overemphasizes the biomedical and ignores psychosocial factors. All of these influences exacerbate and perpetuate the somatic distress of patients with functional somatic syndromes, heighten their fears and pessimistic expectations, prolong their disability, and reinforce their sick role. A six-step strategy for helping patients with functional somatic syndromes is presented here.

Identifiers--Keyword Plus(R): CHRONIC-FATIGUE-SYNDROME;
IRRITABLE-BOWEL-SYNDROME; RANDOMIZED CONTROLLED TRIAL; TERM FOLLOW-UP;
MULTIPLE CHEMICAL SENSITIVITIES; POSTTRAUMATIC-STRESS-DISORDER;
COGNITIVE-BEHAVIORAL THERAPY; MASS PSYCHOGENIC ILLNESS; CARE-SEEKING
BEHAVIOR; UPPER LIMB PAIN

Cited References:

*I MED, 1996, HLTH CONS SERV PERS

9/9/2 (Item 1 from file: 442)
DIALOG(R) File 442:AMA Journals
(c)1999 Amer Med Assn -FARS/DARS apply. All rts. reserv.

00098204
COPYRIGHT American Medical Association 1996

Annual Meeting of American Academy of Otolaryngic Allergy (ARTICLE)

Archives of Otolaryngology
Apr, 1996; Medical News: tzo446
LINE COUNT: 00104
0003-9977

This meeting was held at the Ernest N. Morial Convention Center, New Orleans, La, on September 14 and 16, 1995.

John H. Boyles, Jr, MD, Dayton, Ohio, found in a review of the literature, 65 articles reporting allergy-related inner ear disease. He reported long-term results in 30 personal cases of patients with the classic symptoms of Meniere's disease. Of the patients, 35% had a history of allergies and 53% a family history of Meniere's disease, 47% were sensitive to foods, and 32% were sensitive to chemicals. The workup must include a complete evaluation of inhalants, foods, chemicals, fungal disorders, autoimmune disorders, and nutrition. Eleven of the 30 patients exhibited autoimmune thyroid disorder.

David Morris, MD, La Crosse, Wis, reviewed several double-blind, placebo-controlled studies in southern Europe that demonstrated significant effectiveness of sublingual therapy with grass and house-mite antigens. Recent immunologic studies, which support sublingual treatment, show that pure antigens that reach the bloodstream without adjuvant are, in general, tolerogenic. It is believed that the direct effect on suppressor cells accounts for the positive results in immediate and delayed reactions.

William J. Rea, MD, Richardson, Tex, using flow cytometry, produced DNA histograms showing normal and abnormal cell cycles that indicate immune status. An immune modulator, called **autogenous lymphocyte factor** (ALF), was produced from autogenous cell cultures. The ALF was injected in 101 patients with treatment-failure chemical sensitivity, who had abnormal T cells and T-cell dysfunction. He reported 87% clinical improvement in these patients ($P=.001$). Only six could not tolerate ALF.

Eleven patients were treated with immunotherapy for allergic fungalsinusitis (AFS) by Richard L. Mabry, MD, Scott C. Manning, MD, and Cynthia S. Mabry, RN, Dallas, Tex. They reported on patients whose injections were advanced weekly for an average of 13 months. There were no adverse effects. After 4 to 8 weeks of immunotherapy, crusting was decreased.

William J. Rea, MD, Richardson, Tex, reported autonomic nervous system and immune system abnormalities in patients with silicone breast implants. The patients demonstrated symptoms as follows: musculoskeletal, 87%; neurological, 60%; gastrointestinal, 27%; cardiovascular, 24%; and otologic, 20%. Skin tests for silicone were positive in 68% of patients. Ninety percent of patients demonstrated serologic evidence of autoimmunity. Results of a pupillogram, which reads speed of contraction and dilatation of the pupil as evidence for autonomic dysfunction, was abnormal in 93%.

Andrew J. Lehr, MD, Richard L. Mabry, MD, and Cynthia S. Mabry, RN, Dallas, Tex, demonstrated that a nine-antigen 'midiscreen' was more sensitive than a six-antigen 'miniscreen' and as sensitive as a 13-antigen screening radioallergosorbent test (RAST) to identify patients with significant allergy. The authors concluded that a nine-antigen midiscreen

STIC-ILL

From: Schwadron, Ronald
Sent: Wednesday, February 10, 1999 11:02 AM
To: STIC-ILL
Subject: 08902692

Youdim et al.
Clin. Ecology
vol 7
p 55-61
1990

Youdim et al.
Clin. Ecology
vol 7
p 62-66
1990

Youdim et al.
Clin. Ecology
vol 7
p 67-72
1990

Rea
Annual Meeting-American Academy of Environmental Medicine (I think this is a book of abstracts published by
American academy of Environmental Medicine)
Sept 1995
p213-222

Griffiths et al.
Cytobios
vol 93
p49-66
1998

thnxs

* Please see attached note
from the author.

240959

his phone #

214-368-4132

Env. Health Center of Dallas

s/w
Ellie

x/61

2/10
50

3/1

DIALOG(R)File 65:Inside Conferences

(c) 1999 BLDSC all rts. reserv. All rts. reserv.

01474979 INSIDE CONFERENCE ITEM ID: CN014639864

The Potential Uses for Autogenous Lymphocyte Factors (ALF) in
Environmental Medicine

Rea, William - owner of clinic

CONFERENCE: American Academy of Environmental Medicine: The cutting edge
of environmental medicine-Annual meeting; 30th

ANNUAL MEETING- AMERICAN ACADEMY OF ENVIRONMENTAL MEDICINE, 1995;

30th P: 213-222

The Academy, 1995

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts, presentations,
papers and programme

CONFERENCE SPONSOR: American Academy of Environmental Medicine

CONFERENCE LOCATION: Tucson, AZ

CONFERENCE DATE: Sep 1995 (19950) (19950)

2/3/2

DIALOG(R)File 65:Inside Conferences

(c) 1999 BLDSC all rts. reserv. All rts. reserv.

01474969 INSIDE CONFERENCE ITEM ID: CN014639768

Some Examples of Administrative Law in Action

Rea, W.

CONFERENCE: American Academy of Environmental Medicine: The cutting edge
of environmental medicine-Annual meeting; 30th

ANNUAL MEETING- AMERICAN ACADEMY OF ENVIRONMENTAL MEDICINE, 1995;

30th P: 127-136

The Academy, 1995

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts, presentations,
papers and programme

CONFERENCE SPONSOR: American Academy of Environmental Medicine

CONFERENCE LOCATION: Tucson, AZ

CONFERENCE DATE: Sep 1995 (19950) (19950)



no

520

Handwritten scribbles and signatures.



FAX

Date: 2/23/99

Number of pages including cover sheet: _____

To:
Linda

Phone:

Fax phone: 703/305-3585

CC:

From:
William J. Rea, M.D./Ellie

Phone: (214) 368-4132 373-5161

Fax phone: (214) 691-8432

REMARKS: ☐ Urgent ☐ For your review ☐ Reply ASAP ☐ Please comment

CONFIDENTIALITY NOTICE

The information contained in this facsimile message and accompanying documents is legally privileged and confidential information intended only for the use of the individual or entity named herein. If the reader of this message is not the intended recipient, you are hereby notified that any dissemination, distribution or copy of this telecopy is strictly prohibited. If you have received the telecopy in error, please immediately notify us by telephone (214) 368-4132, and return the original message to us at the address above via the United States Postal Service.

Instructions:

Linda: A paper was not published from the AAEM abstract. However, I am attaching a published paper regarding ALF.

The role of the T lymphocytic cell cycle and an autogenous lymphocytic factor in clinical medicine

Bertie B. Griffiths*, William J. Rea, Bradley Griffiths and Y. Pan

EHC-D Analytical Laboratory, Environmental Health Center - Dallas, 8345 Walnut Hill Lane, Suite 240, Dallas, Texas 75231, U.S.A. (*Reprint address)

Key words: cell cycle, flow cytometry, T lymphocytes, autogenous lymphocytic factor, chemical sensitivity

Abstract

In this study 315 individuals (25 controls, 290 chemically sensitive immuno-compromised patients) were investigated. Each patient had been on a standard therapy of avoidance of pollutants, nutritional supplementation, and injections of antigens for foods, and biological inhalants, but did not attain their immunological competence. Peripheral lymphocytes were collected and DNA histograms were constructed. The flow cytometer was used to evaluate the cell cycle, haematological, and other immunological profiles. From the other portion of the blood specimen, lymphocytes were propagated *in vitro*, harvested, and a lysate, termed the autogenous lymphocytic factor (ALF), was prepared. When treated with ALF, 88% of these individuals showed a significant ($p < 0.001$) clinical improvement which correlated with laboratory findings, involving regulation of abnormal cell cycles, increase in total lymphocytes and subsets T_H , T_G ($p < 0.05$) and cell mediated immunity (CMI) response ($p < 0.001$). The ALF presumably acts as a biological response modifier. The cell cycle and ALF provide clinical tools for diagnosis and regulation of immunological incompetence.

14

Introduction

The cell cycle is the ordered and orderly events of biochemical and morphological sequences, leading from the formation of a daughter cell as a result of mitosis, to the completion of the processes required for its own division into two daughter cells (Lee and Dang, 1995). It is fundamental that following mitosis, two daughter cells are produced. These cells may initiate a new cycle (G_1 phase), some cells may become nonproliferative (G_0 phase) or progress to a restriction point where they are committed to the synthesis of cellular components, (the S phase) and finally complete the cycle (the G_2M phase). The cell cycle then comprises the sum of the growth phases of a specific cell cycle. This cycle is repeated by continuously dividing cells. Even within the same organism the specific cell cycle will vary with different classes of cells. The total time that is necessary to complete the S and G_2 phases is generally constant in different cell types. It seems reasonable then, to assume that most of the time variation takes place in the G_1 phase.

To establish homeostasis, it is imperative that the cell cycle be regulated. To this end, there are biological parameters which may be employed to ascertain the normality of a particular cell cycle. The DNA content in the nucleus of a cell ($2N$ or diploid amount) is constant in all normal organisms (Givan, 1992). There are only two exceptions where the DNA content is not constant; the amount of DNA will vary in cells which undergo meiosis in preparation for sexual reproduction, thus containing the $1N$ or haploid amount of DNA. The other exception applies to those cells which undergo DNA synthesis in preparation for mitosis. These will contain between $2N$ and $4N$ amounts of DNA.

By employing the concept of DNA constancy in a particular organism, and the application of flow cytometric techniques, DNA flow histograms can be constructed depicting a normal cell cycle and the identification of abnormal cycles.

The regulation of the cell cycle in eukaryotes seems to take place at two main transition points, prior to DNA replication at a point in the G_1 phase, termed the restriction point, and prior to cytokinesis at the G_2M phase boundary (Nurse and Bissett, 1981). The progression of the cell cycle from one phase to the other is mediated by specialized proteins, cyclins, and the activation of enzymes called cyclin-dependent kinases (CDKs) and by a number of positive and negative feedback loops. To date seven CDKs have been identified and designated CDK 2 to 8 (Meyerson *et al.*, 1992; Lorinez and Reed, 1984; Nurse, 1994; Sherr, 1994; Helchman and Roberts, 1994; King *et al.*, 1994; Hunter and Pines, 1994). Each CDK acts at different stages of the cell cycle and is differentially regulated by different cyclins (Meyerson and Harlow, 1994). The critical roles played by cyclins in the regulation of eukaryotic cell cycle is amply documented.

These CDKs play very significant roles in the G_1S and G_2M transitions during the mammalian cell cycle. They regulate by phosphorylation, a number of key substrates which subsequently activate a transition from G_1 to S and from G_2 to M (Kamp *et al.*, 1994). The catalytic subunits alone of these CDKs are not active and require the influence of positive regulatory subunits to ensure biochemically active protein kinase holoenzymes (Desai *et al.*, 1992). The positive regulatory subunits employed to this end are cyclins. The activity of CDKs is, therefore, regulated by both cyclins and specific phosphorylation and dephosphorylation (Gold and Nurse, 1989; Draetta and Beach, 1988; Solomon *et al.*, 1990; Gutier *et al.*, 1991; Murray and Krischner, 1989).

Cyclins were identified originally as proteins in the murine invertebrate cells. The concentration of these cyclins accumulate and are destroyed periodically at defined points during the cell cycle (Evans *et al.*, 1983). At present, eight cyclins have been identified; these are designated A, B, C, D, E, F, G₁, and H based on their amino acid sequences (Pines and Hunter, 1989; Xiong *et al.*, 1991), and in some instances, on genetic complementation experiments in yeast (Nurse, 1990; Murray and Krischner, 1989; Lewin, 1990; Sherr, 1993).

The influences of the cyclins are expressed differently; *e.g.* cyclin A exhibits its influence through the S-M phase (Wang *et al.*, 1990; Fang and Newport, 1991; Walker and Maller, 1991; Lehner and O'Farrell, 1990; Pagano *et al.*, 1992; Zindy *et al.*, 1992) of most cells, while cyclin B (which is conserved from yeasts to humans) propels cells into mitosis (Nurse and Bissett, 1981; Evans *et al.*, 1983; Glotzer *et al.*, 1991). Both the A and B cyclins are degraded at the M phase by ubiquitin (Ub)-dependent proteolysis (Glotzer *et al.*, 1991). G₁ cyclins (CIN 1, CIN 2, CIN 3) are assumed to associate with a p34 CDC2 homologue, p34 CDC 28, driving yeast cells into S phase (Reed, 1992; Cross, 1988; Richardson *et al.*, 1989; Hadwiger *et al.*, 1989; Wittenberg *et al.*, 1990; Nasmyth, 1990).

In the Environmental Health Center - Dallas (EHC-D), clinical cases are investigated varying from surgical to environmental illnesses. However, cases involving chemical sensitivity entail the principal aspect of our practice. Over a period of 22 years our laboratory observations show that characteristically, a percentage of the chemically sensitive individuals portray a secondary deficit which is related to immune dysfunction, principally, depression of the suppressor T-cell (Rea *et al.*, 1986, 1987).

Initial profiles of lymphocyte cell cycle progression, T and B lymphocytes, cell mediated immunity and haematograms of patients were compared with those taken subsequent to treatment with autogenous lymphocytic factor (ALF). Acting presumably as a biological response modifier, ALF was observed to modulate deregulated immune profiles.

The focus of this investigation was firstly to observe the patterns of DNA histograms of the T lymphocyte cell cycles, especially in chemically sensitive individuals. The histograms provide a 'snapshot' of the individual's present cell cycle. By comparing an individual's cell cycle with the classic normal cycle, very crucial information will be obtained to enable a scientific regulation of a patient's cell cycle. Secondly the work was to prepare a lysate

from the *in vitro* propagated peripheral T lymphocytes of an individual with an irregular T lymphocytic cell cycle, and observe the regulatory effect of this lysate, when injected subcutaneously into such an individual. Thirdly, the intention was to establish a basis for the regulation of an individual's T lymphocytes which were observed to be irregular due to varied incitants, thus restoring normal T lymphocyte functions, and the enabling of a compromised individual to cope with multiple insults to his/her immune system. Fourthly, we examined the responsiveness of T lymphocytes to autogenous lymphocytic factor (ALF) as measured by the cell mediated immunity (CMI) test, and to establish on an individual basis an index of T lymphocyte ability to respond.

From our search of databases, and to the best of our knowledge there is no documentation pertaining to the regulation of the T lymphocytic cell cycle by lysates of an individual's T lymphocytes, and its application to clinical medicine.

Materials and methods

Isolation of lymphocytes

Venous blood was collected from 315 individuals in heparinized tubes. Of these 25 were healthy employees of EHC-D; these represented normal controls. The other individuals were chemically sensitive, chronically ill patients of the EHC-D and referrals from outside physicians. Erythrocytes were separated by a modified gradient technique using heparinized CPT vacutainers (Becton Dickinson Co.). This preparation was centrifuged for 30 min at 500 x g. The lymphocyte layer (interphase between isolymp and plasma) was removed to sterile polystyrene tubes. Lymphocytic layers were washed three times with normal saline (0.9% sodium chloride) by centrifugation in a refrigerated centrifuge at 500 x g for 20 min. Cells were resuspended in normal saline and counted.

Lymphocyte phenotyping

Monoclonal antibodies against pan T cells, T_{H1} (CD2), T helper cells T_H (CD4), T suppressor/cytotoxic cells T_S (CD8), activated T cells TA1 (CD26) and pan B cells B₄ (CD19) were bought from Coulter Immunology (Hialeah, Florida). Peripheral lymphocytes were obtained by venipuncture, and stained by the use of Coulter Q prep Epics immunology station following manufacturer's instructions.

Preparation of cell cycles

Heparinized whole blood or lymphocytes were used and 100 μ l aliquots of heparinized whole blood or 100 μ l of washed isolated lymphocytes (suspended in normal saline) were placed in 13 x 75 mm tubes. To each tube was added 200 μ l of lysing buffer (Coulter) and gently mixed for ~15 s, then 2.0 ml of DNA stain with RNase was added and mixed for 20 s. The mixture was stained with propidium iodide at a concentration of 50 mg per ml, and allowed a staining period of 15 min and then analysed by flow cytometry.

Flow micro-fluorometric analysis

Cells tagged fluorometrically for DNA content were analysed on the Coulter Epic C flow cytometer, where they were allowed to pass through a laser beam tuned to 488 nm. Fluorescence was measured electronically and recorded as a histogram. DNA distribution in the cell cycle was calculated on accumulated data by parametric analysis to produce a particular DNA histogram.

Preparation of autogenous lymphocytic factor (ALF)**Propagation of lymphocytes**

To cell culture flasks, containing 10 ml RPMI 1640 medium and/or supplemented with 1 ml bovine calf serum was added 1 ml of washed T lymphocytes. The culture was incubated at 37°C and monitored daily for approximately 2 to 3 weeks.

Harvesting of cells

When the cell population approximated 5 to 9 x 10⁶ per ml, culture suspensions were transferred to calibrated 50 ml conical tubes and centrifuged for 30 min at 500 x g. The supernate was discarded, cell peller resuspended in 10 ml normal saline and washed three times by centrifugation for 30 min each time.

Preparation of ALF

Each cell peller was resuspended in ~2.0 ml normal saline and sonicated at 20 W with a duty cycle of 50% for 60 s. The sonicated mixture was sterilized by filtration and stored at -20°C to be diluted for therapeutic use as indicated.

Establishment of a functional T lymphocyte cell cycle

The profile of an individual's T lymphocyte cell cycle may appear normal, yet the lymphocytes may not be functional. A process was established to evaluate the immunological responsiveness of the lymphocytes of the cell cycle.

Stimulation of peripheral lymphocytes

NuncionTM-plastic plates (24 wells) containing 2 ml of RPMI 1640 medium per well were inoculated with ~ 2 to 4×10^6 lymphocytes per ml, 1.25 ng interleukin 1 alpha (IL_1 , alpha) per ml, or with 0.1 ml ALF per ml; negative controls had neither IL_1 nor ALF. The plates were incubated at 37°C for 96 h. Cells were harvested, washed and analysed by flow cytometrically as mentioned above. Comparison of the S and G₂ phases of both the unstimulated and the stimulated T lymphocytes was undertaken and an index computed from the equation:

$$\text{stimulation index} = (\text{stimulated S} + G_2) / (\text{unstimulated S} + G_2)$$

Clinical Investigation

A total of 315 individuals was investigated and of these, 25 were controls. T lymphocyte cell cycle profiles, haematological, T and B profiles, and CMI tests were carried out on all candidates. Individuals with immune deregulation were treated with autogenous lymphocytic factor (ALF). These individuals were chemically sensitive and chronically ill. The illnesses include dermatitis, vasculitis, asthma, organic brain syndrome, and Gulf war syndrome. In all cases, these illnesses were characterized with immune system suppression, dysfunction or deregulation.

Major symptoms presented by these patients include one or more of the following symptoms: lacrimation, pruritus, swelling and puffiness (ocular); fullness, noise, and dizziness (otic); congestion, sneezing, rhinorrhoea and blowing (nasal); lump, clearing and postnasal drip (throat); hypersensitivity reactions (immune); arthritis and arthralgia, fatigue and muscle pain (musculoskeletal); pressure and cough (chest); weight loss and fatigue (constitutional); miliary, ethmoidal and frontal (headache); insomnia, shortness of breath and depression (neurological).

All 290 chemically sensitive individuals presented a history of being affected by environmental incitants found in categories such as food, biological inhalants, and chemicals. They presented histories of varied backgrounds, but common among them was that all showed irregular cell cycles and abnormal T lymphocyte profiles in both numbers and functions in T and B lymphocytes and subsets. DNA histograms showed over or under accumulation of various subtypes of lymphocytes in one or more phases of the cell cycle of each individual.

Cell-mediated Immunity

Delayed cutaneous hypersensitivity, or cell-mediated immunity (CMI) responses were evaluated in 190 patients; the results were recorded before and after treatment with ALF. The multitest CMI test kit (Mérieux Institute, Miami, Florida) was used. Each kit contained seven antigens: tetanus, diphtheria, candida, proteus, streptococcus, trichophyton, and tuberculin. The tests were evaluated and read at 48 h. Evaluation involves scoring the size, and number of increase of the wheals. The diameter of each induration was measured in millimetres and averaged. A reaction was considered positive if the average diameter was 2 mm or more.

Comparative evaluation of T and B lymphocytes

T and B lymphocytes and subsets were evaluated flow cytometrically before and after treatment with ALF.

Table 1 Representative profile of symptoms and signs after autogenous lymphocytic factor treatment

Symptoms	Improvement:		No change:		Deterioration:		Total patients
	Number	%	Number	%	Number	%	
Hypersensitive reaction	63	63	33	33	1	1	100
Recurrent infection	38	57	29	43	0	0	67
Fatigue	60	68	27	31	1	1	88
Lack of concentration	43	54	36	46	0	0	79
Arthritis	19	44	23	54	1	2	43
Gastrointestinal upset	29	40	43	60	0	0	72
Headache	28	44	33	53	2	3	64
Depression	42	58	30	42	0	0	72

Establishment of therapeutic dosage of ALF

In vitro and *in vivo* procedures were used to establish dosage. Blastogenesis was the *in vitro* test employed for establishing dosage in general. *In vitro* grown peripheral lymphocytes were challenged comparatively with the known mitogens phytohaemagglutinin (PHA), concanavalin A (Con A), and pokeweed mitogen (PWM) of varied concentrations. A dose responsive curve was established to evaluate the efficacy of ALF with that of the mitogens; 0.1 ml. of 1:10 dilution of ALF was found to be optimal when administered twice weekly. Intradermal test was carried out on all recipients depending on their response and sensitivity. If unusual sensitivities were observed, the ALF was further diluted. This procedure was successively repeated until no adverse reaction was observed.

Results

Significant changes were observed within 1 to 6 weeks in patients treated with ALF. Changes were observed in immune regulation and overall clinical manifestations. With regard to clinical manifestations, there were noteworthy improvements (Table 1), although minimal symptoms continued after approximately 3 weeks of continued therapy.

In a normal cell cycle, the highest percentage of lymphocytes should be in the G_0 to G_1 phase (Figure 1). This percentage will change dramatically when these lymphocytes are stimulated by various incitants (Figures 3a and 3b). Consequently, different percentages will appear in the S and G_2 M phases, producing a deregulated profile (Figures 1 to 4). Treatment with ALF regulates the T lymphocyte cell cycle profile.

Immunologically, there were significant regulations of T lymphocyte cell cycles, especially from one phase of the cycle to another. Patients became less sensitive and more tolerant to specific incitants. As treatment continued, in general in about 6 weeks, a more drastic shift toward that of a normal profile was observed. Figures 1 to 4 summarize the regulatory changes of some of the cell cycles studied.

Changes were observed in the profiles of the T and B cells where T and B lymphocytes and their subsets were evaluated before and after treatment. There was a significant ($p < 0.01$) change in the total lymphocyte count and subsets, T_4 , T_8 in 92 patients investigated. It should be noted that ALF seems to act as an immune modifier since the total lymphocytes, T_4 and T_8 were significantly elevated or reduced in order to maintain normalization (Tables 2 and 3).




Figure 1 Diagrammatic representation of a normal mammalian cell cycle, showing the relative number of lymphocytes in each phase of the cycle, as displayed by the intensity of fluorescence of the fluorochrome-bound DNA in each lymphocyte. Normally, the percentage of lymphocytes in each phase of the cell cycle is 90-95, 5-10 and 0-5 in the G_0G_1 phase, S phase and G_2M phase, respectively. These quiescent lymphocytes display, normally, a low level of replication, and progression through the cell cycle. In response to stimuli, these cells undergo rapid division and cycle rapidly. The shadings show that there is overlapping of phases and not specific lines of demarcation: stippled, G_0G_1 phase (90-95% noncycling lymphocytes); hatched, S phase (5-10% cycling lymphocytes); unshaded (open), G_2M phase (0-5% cycling lymphocytes); and solid areas indicate overlapping of phases.

Table 2 Representative changes in T and B cell profiles after 92 chemically sensitive patients were treated with autogenous lymphocytic factor

Treatment	Total lymphocytes	T_{11}	T_4	T_8	T_4/T_8	B_4
Before	2,112 \pm 692	1,624 \pm 457	930 \pm 45	439 \pm 58	2.3 \pm 0.8	188 \pm 102
After	2,232 \pm 678*	1,634 \pm 544	1,027 \pm 297**	478 \pm 189**	1.3 \pm 7.4	171 \pm 120

* $p < 0.05$; ** $p < 0.01$; $n = 92$.




Figure 2 A representative DNA histogram of peripheral T lymphocytes from normal volunteers. Note that the highest percentage of lymphocytes is in the G_0G_1 phase, as expected, but the percentage varies in different individuals in keeping with the prevalence of environmental pollutants. Thus, the ideal normal profile is seldom achieved. The abscissa shows the relative fluorescence of the fluorochrome-bound DNA in each cell.

Table 3 Profile of T lymphocyte subsets modulation after treatment with autogenous lymphocytic factor

After treatment	Total lymphocytes:		T_{H1}		T_H		T_H	
	Number	%	Number	%	Number	%	Number	%
Increase	52	57	41	46	53	53	55	60
Decrease	40	43	51	54	39	42	37	40
Probability	<0.05		>0.05		<0.05		<0.01	

Table 4a Typical cell-mediated response in (number and size) chemically sensitive individuals treated with autogenous lymphocytic factor

Patients	Before ALF	After ALF	P
190	5.46 \pm 5.81	9.28 \pm 7.25	<0.001

Results are given as mean \pm SD.

Patients showed significant improvement ($p < 0.001$) in their CMI scores (Table 4). The regulatory effect of the immune system can be objectively assessed periodically after the initial treatment with ALF. The resourceful parameters are profiles of T lymphocyte cell cycle, T and B lymphocytes, and their subsets, cell-mediated immunity, signs and symptoms.

Side effects of ALF

The side effects were minimal and occurred only in six adults (five females and one male) where the average age was 52 years. These patients were intolerant of ALF, their symptoms included pain and irritation in the throat, burning in the eyes, pain and irritation in the chest, heat palpitations, influenza-like symptoms, headache, fatigue, and chills.

Table 4b Cell-mediated immunity positive score (number and size) after treatment with autogenous lymphocytic factor

Score	Patients	%	P
Increase	58	74	<0.001
No change	9	12	
Decrease	11	14	




Figure 3a A representative irregular T lymphocyte cell cycle profile showing the effect(s) of stimulating the environmentally ill patients influence, primarily, the lymphocytes in the S phase.

Figure 3b A representative irregular T lymphocyte cell cycle profile showing the effect(s) of the stimulating incitants on the lymphocytes in the G₂M phase. It seems that each incitant, or a mixture of incitants affects lymphocytes in a particular phase(s) of the cell cycle, resulting in a variety of irregular cell cycle profiles and presumably dictates varied patterns of clinical manifestations. The abscissae in Figures 3a and 3b show the relative fluorescence of the fluorochrome-bound DNA in each cell.

Discussion

The flow cytometric profile of the T lymphocyte cell cycle as demonstrated by DNA histogram presents a reflection of the status of T lymphocytes in an individual. Of great importance is that its application is not limited to a certain category of individuals, but to normal subjects as well as individuals who are compromised by varied incitants. The essence of its clinico-biological importance, as detected in the present investigation, is that it is a reflection of T lymphocyte cell function, and may facilitate an in-depth approach to the treatment of some immunodeficient illnesses.

The progression of T lymphocytes from one phase of the cell cycle to another is time-dependent, ~8 to 12 h in the G₀-G₁ phase, 6 to 8 h in the S phase, and 0.5 to 1 h in the G₂M phase. The 290 individuals who were investigated in this study were affected principally by environmental incitants in categories such as foods, biological inhalants, and chemicals. They presented histories of varied backgrounds and different cell cycle profiles. The DNA histogram




Figure 4a Initial T lymphocyte cell cycle profile of a patient on examination at the clinic. Note that the irregularity of the profile is emphasized in the G₂M phase.

Figure 4b Cell cycle of the patient in Figure 4a after the subject was treated with autogenous lymphocytic factor (ALF) for 3 weeks. Note that the percentages of lymphocytes in each phase of the cell cycle are still irregular, but there has been a significant improvement. The highest percentage of lymphocytes now appears in the G₀G₁ phase.

Figure 4c T lymphocyte cell cycle of patient (Figures 4a and b) subsequent to 3 months treatment with ALF. Note that the cell cycle has attained a relatively normal profile. ALF seems to act as a biological response modifier. The abscissae in Figures 4a, b and c show the relative fluorescence of the fluorochrome-bound DNA in each cell.

cycles showed over or under accumulation of lymphocytes in one or more phases of the cell cycle of each individual (Figures 3a and 3b). As shown in Figure 2, even the normal controls do not present an ideal DNA histogram profile of T lymphocytes. This is due to the fact that, in general, the ideal environment is seldom ever achieved, thus there is always some degree of immunological compromise to most individuals. It is reasonable then to assume that the incitant(s) which lead to immuno-incompetence of these individuals are capable of inactivating the particular enzyme(s), cyclin-dependent kinases, or depleting specific cyclins whose combination with their specific kinases are instrumental in catalysing the progression of T lymphocytes from one phase to another of the cell cycle. The T lymphocytes of the individual affected are locked in a particular phase, either resting, synthesizing, or multiplying too much in the G_0 - G_1 , S, G_2 M phase respectively. Thus the subject manifests symptoms peculiar to the phases(s) affected. This hypothesis offers an opportunity to associate clinical manifestations with T lymphocyte cell cycle irregularity.

The treatment of lasting importance would be reasonably thought of as a biological response modifier, which would stimulate CDKs, regulate the cell cycle and the enzymes of purine and pyrimidine nucleotide synthesis. It is now generally accepted that these enzymes are elevated during the S phase (Cory, 1993). It seems logical that ALF stimulates these regulatory functions. The autogenous lymphocytic factor is a dialysable mixture of the many effector substances which may be released from *in vitro* grown stimulated lymphocytes with the ability to invoke immunological influences *in vivo* or *in vitro*, (Griffiths and Rea, 1995). The expected biological activity(s) of ALF would be to act as a biological response modifier, with mechanisms of action as suppressor and regulator of the immune system, especially in the regulation of the T lymphocyte cell cycle.

Kamp *et al.* (1994) showed that progression through the cell cycle requires the joint influence of positive regulator subunits, cyclins, and cyclin dependent kinases. These subunits regulate by phosphorylation a number of key substrates which subsequently activate a transition from G_1 to S, and G_2 to M phases of the cell cycle. Phosphate groups are transferred from ATP to a special amino acid in the target protein by protein kinases, while phosphatases remove the phosphate groups from the target proteins. The addition and removal of phosphate groups significantly affect the biochemical behaviour of the target proteins. Many protein kinases and phosphatases have a specific affinity for their target proteins, and act as determinants for controlling the activity(s) of their target

proteins. Indeed, ALF may possess these protein kinases which, acting as molecular switches, regulate an irregular T lymphocyte cell cycle to that of an ordered and orderly progression.

As demonstrated (Figure 4c), ALF regulates deregulated cell cycles, improves immunological profiles by increasing or decreasing the number of circulating lymphocytes and their subpopulations (Tables 2 and 3); and restores immune responses as demonstrated by enhanced cutaneous hypersensitivity (CMI) (Table 4). ALF also invoked immediate intradermal test response within 45 min of administration subsequent to the administration of antigen challenges which were previously negative or delayed for at least 15 days. Responses were observed by the number and size of wheals to the antigens which were inoculated. It is noteworthy that some chemically sensitive patients responded favourably, or completely recovered after environmental exposure to air, food, water, nutritional support and exercise. Some patients responded but did not fully recover. The patients who recovered reacted profoundly to exposure of very minute environmental insults of ambient chemicals, biological inhalants, and foods, or inhalants. However, when these patients were treated with ALF, their hypersensitivity was markedly reduced or disappeared; there was also significant improvement in recurrent infection, fatigue, headaches, depression, concentration, and gastrointestinal upsets. Eight patients immediately improved on initial injection with ALF.

T-suppressor, T-helper cells, and total lymphocytes were increased as reflected in an increase in a low population, and a decrease in a too high population. This suggests that ALF is a modulator rather than a stimulator. The mechanism(s) of action of ALF have not yet been ascertained. Indeed, the purification and determination of ALF from sorted T lymphocytes with or without helper phenotypes will be necessary to facilitate understanding the mechanism of action. However, it has been documented that the human cells contain a regulatory protein, CKS protein, which is a genetic suppressor of temperature sensitive CDK mutations. There are two isoforms of this CKS protein, namely CKSH1 and CKSH2. It is believed that CKSH2 protein binds to the catalytic subunit of the CDKs, and is essential for their biological function (Parge *et al.*, 1993). To this end, further investigations are being carried out.

The concept of a functional cell cycle is timely. Indeed, the profile of a cell cycle offers an insight into the immune status of a patient. However, the cell cycle profile does not indicate whether or not the lymphocytes are immunologically responsive or capable of responding to therapy. The influence of varied environmental incitants and/or

other systemic illnesses may render these lymphocytes incapable of normal progression through the cell cycle, thus resulting in a non-functional cell cycle. *In vitro* lymphocyte activation represents a standard procedure for evaluating cell mediated responses to a variety of stimuli including antibodies, polyclonal mitogens, specific antigens and cytokines. Interleukin-1 alpha (IL₁ alpha) and ALF were used as activators in the present investigation. Interleukin-1 alpha was used due to its extremely broad spectrum of bioactivity. ALF was used to observe its capability as a biological response modifier. The stimulatory process presented in this investigation offers a method of computing lymphocyte stimulatory response, and consequently an opportunity to establish an individual regime of treatment of illnesses associated with immunodeficiency. This investigation offers: (1) a vivid picture of the status of the immune profile of a patient as demonstrated by the DNA histogram of the T lymphocyte cell cycle; (2) an opportunity to regulate an irregular T lymphocyte cell cycle by treatment with autogenous lymphocytic factor; and (3) a clinical tool for the regulation of abnormal cell cycles and/or haematological and immunological profiles in humans with immunological deregulation. It should be noted that there is a pending United States patent, number 08/380,063 for the application of the cell cycle to clinical medicine, and also the regulation of the cycle by autogenous lymphocytic factor.

Acknowledgements

The authors are grateful to Dr Alfred Johnson and Dr Gerald Ross of the Environmental Health Center - Dallas, for their contribution to the clinical data, and to Dr Vernon E. Scholes for reviewing this manuscript.

References

- CORY J. G. 1993. Purine and pyrimidine nucleotide metabolism. In *Biochemistry with Clinical Correlation*. pp 529-73. Edited by T. M. Devlin. Wiley-Liss, New York.
- CROSS F. R. 1988. DAF 1 a mutant affecting size control, pheromone arrest, and cell kinetics of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 8 4675-84.
- DESAI D., Gu Y. and Morgan D. O. 1992. Activation of human cyclin-dependent kinases *in vitro*. *Mol. Cell. Biol.* 3 671-82.
- DRAETCA G. and Beach D. 1988. Activation of CDC2 protein kinase during mitosis in human cells: cell cycle-dependent phosphorylation and subunit rearrangement. *Cell* 3 17-26.
- EVANS T. E., Rosenthal E. T., Youngblom J., Distel D. and Hunt T. 1983. Cyclin: a protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell* 33 389-98.
- FANG F. and Newport J. W. 1991. Evidence that the G₁-S and G₂-M transitions are controlled by different CDC2 proteins in higher eukaryotes. *Cell* 68 731-42.
- GIVAN A. L. 1992. Cell from within. In *Flow Cytometry - First Principles*. pp 103-33. Edited by A. L. Givan. Wiley-Liss, New York.

- GLOTZER M., Murray W. A. and Kirschner M. W. 1991. Cyclin is degraded by the ubiquitin pathway. *Nature* 349 132-8.
- GOLD K. L. and Nurse P. 1989. Tyrosine phosphorylation of the fission yeast CDC2+ protein kinase regulated entry into mitosis. *Nature* 324 38-45.
- GRIFFITHS B. B. and Rea W. J. 1995. Pending U.S. patent number 08/380,063.
- GUTIER J., Solomon M. J., Bocher R. N., Bazan J. F. and Kirschner M. W. 1991. CDC25 is a specific tyrosine phosphatase that directly activates p34 CDC2. *Cell* 67 197-211.
- HADWIGER J. A., Wittenberg C., Richardson H. E., Lopes M. and Reed S. I. 1989. A family of cyclin homologs that control the G phase in yeast. *Proc. natn. Acad. Sci. U.S.A.* 86 6255-9.
- HELMAN K. A. and Roberts J. M. 1994. Rules to replicate by. *Cell* 78 567-82.
- HUNTER T. and Pines J. 1994. Cyclins and cancer 11: cyclin D and CDK inhibitors come to age. *Cell* 79 573-82.
- KAMP A., Gruis N. A., Weaver-Feldhaus J., Liu Q., Harshman K., Tavtigian S. V., Stockert E., Day R. S. III, Johnson B. E. and Skolnick M. J. 1994. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 264 436-40.
- KING R. W., Jackson P. K. and Kirschner M. W. 1994. Mitosis in transition. *Cell* 79 557-62.
- LEE W. M. and Dang C. V. 1995. Control of cell growth, differentiation and death. In *Hematology, Basic Principles and Practice*. pp 69-84. Edited by R. Hoffman, E. KaBanz Jr, S. J. Shattil, B. Furie, H. J. Cohen and L. E. Silberman. Churchill Livingstone, New York.
- LEHNER C. and O'Farrell P. 1990. The rule of Drusophilia cyclin A and B in mitotic control. *Cell* 61 535-47.
- LEWIN B. 1990. Driving the cell cycle: M phase kinase, its partners, and substrates. *Cell* 61 743-52.
- LORINEZ A. T. and Reed S. I. 1994. Primary structure homology between the product of yeast cell division control gene CDC-28 and vertebrate oncogenes. *Nature* 307 183-5.
- MEYERSON M. and Harlow E. 1994. Identification of G₁ kinase activity for CDK6, a novel cyclin D partner. *Mol. Cell. Biol.* 14 2077-86.
- MEYERSON M., Enders G. H., Wia C. L., Su L., Gorka C., Nelson C., Harlow E. and Tsai L. 1992. A family of human CDC2-related protein kinases. *EMBO J.* 11 2909-17.
- MURRAY A. W. and Kirschner M. W. 1989. Cyclin synthesis drives the early embryonic cell cycle. *Nature* 339 275-80.
- MURRAY A. W. and Kirschner M. W. 1989. Dominoes and clocks: the union of two views of the cell cycle. *Science* 246 614-21.
- NASMYTH K. A. 1990. Far-reaching discoveries about the regulation of START. *Cell* 63 1117-20.
- NURSE P. and Bissett Y. 1981. Gene required in G₁ for commitment to cell cycle and in G₂ for control of mitosis in fission yeast. *Nature* 292 558-60.
- NURSE P. 1990. Universal control mechanism regulating onset of M-phase. *Nature* 344 503-8.
- NURSE P. 1994. Ordering S phase and M phase in the cell cycle. *Cell* 79 547-50.
- PAGANO M., Pepperkok P., Verde F., Ansorge W. and Draetta G. 1992. Cyclin A is required at two points in the human cell cycle. *EMBO J.* 11 961-71.
- PARGE H. E., Arvai A. S., Murtari D. T., Reed S. I. and Tainer J. A. 1993. Human CksHs2 atomic structure: a role for its hexameric assembly in cell cycle control. *Science* 262 387-95.
- PINES J. and Hunter T. 1989. Isolation of human cyclin C DNA: evidence for cyclin M-RNA and protein regulation in the cell cycle and for interaction with P34 CDC2. *Cell* 58 833-46.
- REA W. J., Pan Y., Johnson A. R., Fenyves E. J. and Smadl N. 1986. T and B lymphocyte parameters measured in chemically sensitive patients and controls. *Clin. Ecol.* 4 11-4.
- REA W. J., Pan Y., Johnson A. R. and Fenyves E. J. 1987. T and B lymphocytes in chemically sensitive patients with toxic volatile organic hydrocarbons in their blood. *Clin. Ecol.* 5 171-5.
- REED S. I. 1992. G₁ specific cyclins in search of an S-phase-promoting factor. *Trends Genet.* 7 95-9.

- RICHARDSON H. E., Wittenberg C., Cross F. and Reed S. I. 1989. An essential G₁ function for cyclin-like proteins in yeast. *Cell* 59 1127-33.
- SHERR C. J. 1993. Mammalian G₁ cyclins. *Cell* 73 1059-65.
- SHERR C. J. 1994. G₁ phase progression: cycling on cue. *Cell* 79 551-5.
- SOLOMON M. J., Glotzer M., Lee T. H., Philippe M. and Krischner M. W. 1990. Cyclin activation of p34 CDC2. *Cell* 63 1013-24.
- WALKER D. H. and Maller J. L. 1991. Role of cyclin A in the dependence of mitosis on completion of DNA replication. *Nature* 354 314-7.
- WANG J., Chenevasse X., Hanglein B. and Brechot C. 1990. Hepatitis B virus Integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 343 555-7.
- WITTENBERG C., Sugimoto K. and Reed S. I. 1990. G₁-specific cyclins of *S. cerevisiae*: cell cycle periodicity, regulation by mating pheromone, and association with the p34 CDC28 protein kinase. *Cell* 62 225-37.
- XIONG Y., Connolly T., Futcher B. and Beach D. 1991. Human D-type cyclin. *Cell* 65 691-8.
- ZINDY F., Lamas E., Chenevasse X., Sobezak J., Wang J., Fesquet D., Hanglein B. and Brechot C. 1992. Cyclin A is required in S phase in normal epithelial cells. *Biochem. biophys. Res. Commun.* 182 1144-54.

Accepted 20 January 1998

News to authors

Manuscripts will be refereed, processed and published rapidly providing the typescript and illustrations have been carefully and accurately prepared in the correct style of each journal.

By following the style of our biomedical journals meticulously you can obtain the advantages of some of the most rapid publication rates for research papers available anywhere. However, a prerequisite is that manuscripts must be impeccably presented in the journal style.

Read the leaflets prepared for authors, entitled *Information for contributors* and *Photographic illustrations*, and send manuscripts for the international biomedical journals MICROBIOS, CYTOBIOS and *BIOMEDICAL LETTERS*, to Dr Stuart Anderson, Executive Editor, The Faculty Press, 88 Regent Street, Cambridge CB2 1DP, England.

- * MICROBIOS is a biomedical research journal, established in 1969, which is concerned with all aspects of Bacteriology and Microbiology. Issues are published every three to four weeks comprising four volumes per annum.
- * *BIOMEDICAL LETTERS* is an international journal for rapid publication of medical, biomedical, and neuroscience research papers, and was first published in 1976. Issues are despatched bimonthly.
- * CYTOBIOS was founded in 1969, and is a biomedical journal for research papers into all aspects of cell science and genetics. Issues are published monthly in four volumes per annum.
- * Manuscripts are peer reviewed.
- * Fifty reprints are provided free to the first named author, although postage is extra.
- * Worldwide distribution, so authors invariably receive many requests for reprints.
- * Abstracted in CURRENT CONTENTS and all the leading abstracting journals.
- * Subscription rates and leaflets for authors are available from the publishers.

The Faculty Press 88 Regent Street Cambridge CB2 1DP England

MICROBIOS

is an international biomedical research journal, established in 1969, which is devoted to fundamental studies of viruses, bacteria, microfungi, microscopic algae, and protozoa. It is concerned with all aspects of micro-organisms, but lays particular emphasis upon chemical microbiology.

Original observations are accepted on the applications of microbiology in the fields of pharmaceutical and chemical production; food manufacture and spoilage; public health and sanitation; biodeterioration; pharmacology and immunology.

Papers on the organization and metabolic activities of micro-organisms are published, as well as work on cell-virus interactions. Manuscripts which are especially welcome are those dealing with the chemical anatomy of micro-organisms, and the biochemical and biophysical factors that affect microbial activity.

The subscription rate for 1998 will be £395.00 sterling.

CYTOBIOS

is a transworld biomedical research journal, established in 1969, which publishes original investigations into all aspects of cell organization. Contributions will be accepted on the behaviour, structure and function of animal and plant cells, including studies on extracellular products and subcellular organelles.

The journal emphasizes work at chemical and molecular levels. It publishes original papers on cytogenetics; cell division and growth; cell physiology and pathology; immunochemistry and immunobiology. Manuscripts are solicited which correlate findings in the biochemical and biophysical fields with morphological, cytological and physiological knowledge.

Discoveries resulting from advances in, and application of, modern biological and medical techniques to cytology are particularly welcome. So also are cytochemical papers which contribute to an understanding of cell organization and to the study of organic fine structure.

The subscription rate for 1998 will be £395.00 sterling.

BIOMEDICAL LETTERS

is an international research journal, established in 1976 as *Microbios Letters*, having the fundamental aim of accelerated publication, and distribution to a worldwide readership. It is intended for short and preliminary biomedical communications, but may include some longer papers and reviews. In general manuscripts should not exceed 5,000 words in length and include only one or two Tables and/or Figures.

BIOMEDICAL LETTERS is primarily designed for the publication of medical research papers. Clinical studies will be considered, and papers in such fields as cellular pharmacology, virology, bacteriology, biochemistry, immunology, molecular biology, biochemical genetics, biophysics, haematology, physiology. Manuscripts on neuroscience, radiation biology and cancer research, will be particularly welcome.

The subscription rate for 1998 will be £200.00 sterling.

Reply slip

Manuscripts

To enable the Executive Editors to plan the publishing programme of forthcoming issues, the following information will be much appreciated:

- | | |
|------------------------------------------------------------------------------------------|--------------------------|
| (1) I hope to submit a paper for publication in | Tick box |
| (a) MICROBIOS | <input type="checkbox"/> |
| (b) CYTOBIOS | <input type="checkbox"/> |
| (c) BIOMEDICAL LETTERS | <input type="checkbox"/> |
| (2) Please send me a free copy of the leaflet
entitled 'Information for contributors' | <input type="checkbox"/> |

The probable title of the paper will be:

.....

.....

.....

.....

I confirm that this manuscript will be based on original, unpublished research, and I understand that all papers are subject to peer reviewing procedures before acceptance. The approximate date of submission will be:

.....

Name

Status

Address



.....

.....

.....

Please complete and return to the address below:

THE FACULTY PRESS 88 Regent Street Cambridge CB2 1DP Great Britain



**American
Academy of
Environmental
Medicine**

For more information, please write:

American Academy of Environmental Medicine
Box CN 1001-2001
New Hope, PA 18938

Phone: (215) 862-4544

Fax: (215) 862-4583

Email: aaem@bellatlantic.net

All contributions are tax deductible.





Creation date: 02-28-2005
Indexing Officer: TBERHE1 - TERHASE BERHE
Team: OIPEScanning
Dossier: 09771346

Legal Date: 01-26-2001

No.	Doccode	Number of pages
1	IIFW	1
2	SRFW	1
3	FWCLM	1
4	TRNA	2
5	DRW	4
6	SPEC	15
7	CLM	1
8	ABST	1
9	OATH	3
10	BIB	1
11	WFEE	1

Total number of pages: 31

Remarks:

Order of re-scan issued on